

# SGPP1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13228a

## Product Information

---

<b>Application</b>	IHC-P-Leica, WB, E
<b>Primary Accession</b>	<a href="#">Q9BX95</a>
<b>Other Accession</b>	<a href="#">NP_110418.1</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB33048
<b>Calculated MW</b>	49108
<b>Antigen Region</b>	13-42

## Additional Information

---

<b>Gene ID</b>	81537
<b>Other Names</b>	Sphingosine-1-phosphate phosphatase 1, SPPase1, Spp1, hSPP1, hSPPase1, 313-, Sphingosine-1-phosphatase 1, SGPP1
<b>Target/Specificity</b>	This SGPP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 13-42 amino acids from the N-terminal region of human SGPP1.
<b>Dilution</b>	IHC-P-Leica~~1:500 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	SGPP1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	SGPP1 ( <a href="#">HGNC:17720</a> )
<b>Function</b>	Specifically dephosphorylates sphingosine 1-phosphate (S1P), dihydro-S1P, and phyto-S1P. Does not act on ceramide 1-phosphate, lysophosphatidic acid

or phosphatidic acid (PubMed:[16782891](#)). Sphingosine-1-phosphate phosphatase activity is needed for efficient recycling of sphingosine into the sphingolipid synthesis pathway (PubMed:[11756451](#), PubMed:[12815058](#), PubMed:[16782891](#)). Regulates the intracellular levels of the bioactive sphingolipid metabolite S1P that regulates diverse biological processes acting both as an extracellular receptor ligand or as an intracellular second messenger (PubMed:[11756451](#), PubMed:[12815058](#), PubMed:[16782891](#)). Involved in efficient ceramide synthesis from exogenous sphingoid bases. Converts S1P to sphingosine, which is readily metabolized to ceramide via ceramide synthase. In concert with sphingosine kinase 2 (SphK2), recycles sphingosine into ceramide through a phosphorylation/dephosphorylation cycle (By similarity). Regulates endoplasmic-to-Golgi trafficking of ceramides, resulting in the regulation of ceramide levels in the endoplasmic reticulum, preferentially long-chain ceramide species, and influences the anterograde membrane transport of both ceramide and proteins from the endoplasmic reticulum to the Golgi apparatus (PubMed:[16782891](#)). The modulation of intracellular ceramide levels in turn regulates apoptosis (By similarity). Via S1P levels, modulates resting tone, intracellular Ca(2+) and myogenic vasoconstriction in resistance arteries (PubMed:[18583713](#)). Also involved in unfolded protein response (UPR) and ER stress-induced autophagy via regulation of intracellular S1P levels (PubMed:[18583713](#), PubMed:[20798685](#)). Involved in the regulation of epidermal homeostasis and keratinocyte differentiation (By similarity).

<b>Cellular Location</b>	Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell membrane {ECO:0000250 UniProtKB:Q9J199}; Multi-pass membrane protein
<b>Tissue Location</b>	Ubiquitous, with the strongest level in placenta and kidney.

## Background

---

Sphingosine-1-phosphate (S1P) is a bioactive sphingolipid metabolite that regulates diverse biologic processes. SGPP1 catalyzes the degradation of S1P via salvage and recycling of sphingosine into long-chain ceramides (Mandala et al., 2000 [PubMed 10859351]; Le Stunff et al., 2007 [PubMed 17895250]).[supplied by OMIM].

## References



---

Hicks, A.A., et al. PLoS Genet. 5 (10), E1000672 (2009) :  
 Le Stunff, H., et al. J. Biol. Chem. 282(47):34372-34380(2007)  
 Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)  
 Olsen, J.V., et al. Cell 127(3):635-648(2006)  
 Giussani, P., et al. Mol. Cell. Biol. 26(13):5055-5069(2006)


## Images

---

Anti-SGPP1 Antibody (N-term) at 1:1000 dilution + human placenta lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 49 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



SGPP1 Antibody (N-term) (Cat. #AP13228a) western blot analysis in NCI-H292 cell line lysates (35ug/lane). This demonstrates the SGPP1 antibody detected the SGPP1 protein (arrow).



SGPP1 Antibody (N-term) (Cat. #AP13228a) immunohistochemistry analysis in formalin fixed and paraffin embedded human placenta tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of SGPP1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.