

SEC14L4 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13245a

Product Information

| | |
|--------------------------|-----------------------------|
| Application | WB, IHC-P, E |
| Primary Accession | Q9UDX3 |
| Other Accession | NP_777637.1 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB33117 |
| Calculated MW | 46644 |
| Antigen Region | 41-70 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 284904 |
| Other Names | SEC14-like protein 4, Tocopherol-associated protein 3, SEC14L4, TAP3 |
| Target/Specificity | This SEC14L4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 41-70 amino acids from the N-terminal region of human SEC14L4. |
| Dilution | WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | SEC14L4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-----------------|--|
| Name | SEC14L4 |
| Synonyms | TAP3 |
| Function | Probable hydrophobic ligand-binding protein; may play a role in the transport of hydrophobic ligands like tocopherol, squalene and |

phospholipids.


Background

The protein encoded by this gene is highly similar to the protein encoded by the *Saccharomyces cerevisiae* SEC14 gene. The SEC14 protein is a phosphatidylinositol transfer protein that is essential for biogenesis of Golgi-derived transport vesicles, and thus is required for the export of yeast secretory proteins from the Golgi complex. The specific function of this protein has not yet been determined. Alternative splicing results in multiple transcript variants.


References

Mokashi, V., et al. *Biochem. Biophys. Res. Commun.* 316(3):688-692(2004)
Collins, J.E., et al. *Genome Biol.* 5 (10), R84 (2004) :
Kempna, P., et al. *Free Radic. Biol. Med.* 34(11):1458-1472(2003)
Dunham, I., et al. *Nature* 402(6761):489-495(1999)

Images



SEC14L4 Antibody (N-term) (Cat. #AP13245a) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the SEC14L4 antibody detected the SEC14L4 protein (arrow).



SEC14L4 Antibody (N-term) (Cat. #AP13245a) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of SEC14L4 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.