10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# OR4X2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13266b

## Product Information

| | |
|---|---|
| **Application** | WB, E |
| **Primary Accession** | Q8NGF9 |
| **Other Accession** | NP_001004727.1 |
| **Reactivity** | Human |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB33177 |
| **Calculated MW** | 34289 |
| **Antigen Region** | 276-303 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 119764 |
| **Other Names** | Olfactory receptor 4X2, Olfactory receptor OR11-105, OR4X2 |
| **Target/Specificity** | This OR4X2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 276-303 amino acids from the C-terminal region of human OR4X2. |
| **Dilution** | WB~~1:1000 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | OR4X2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | OR4X2 |
| **Function** | Odorant receptor. |
| **Cellular Location** | Cell membrane; Multi-pass membrane protein. |


## Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

## References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004) Fuchs, T., et al. Genomics 80(3):295-302(2002)

## Images




OR4X2 Antibody (C-term) (Cat. #AP13266b) western blot analysis in human normal Uterus tissue lysates (35ug/lane).This demonstrates the OR4X2 antibody detected the OR4X2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.