

TRIM54 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13292a

Product Information

Application WB, IHC-P, E Primary Accession Q9BYV2

Other Accession <u>Q58D15, NP 912730.2, NP 115935.3</u>

Reactivity Human **Predicted** Bovine Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB33259 40301 **Calculated MW Antigen Region** 1-30

Additional Information

Gene ID 57159

Other Names Tripartite motif-containing protein 54, Muscle-specific RING finger protein,

MuRF, Muscle-specific RING finger protein 3, MuRF-3, MuRF3, RING finger

protein 30, TRIM54, MURF, MURF3, RNF30

Target/SpecificityThis TRIM54 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 1-30 amino acids from the N-terminal

region of human TRIM54.

Dilution WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions TRIM54 Antibody (N-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name TRIM54

Synonyms MURF, MURF3, RNF30

Function May bind and stabilize microtubules during myotubes formation.

Cellular Location Cytoplasm, cytoskeleton. Cytoplasm, myofibril, sarcomere, Z line.

Note=Associates with microtubules. Localizes to the Z-lines in skeletal

muscles (By similarity).

Tissue Location Specifically expressed in heart and skeletal muscle.


Background

The protein encoded by this gene contains a RING finger motif and is highly similar to the ring finger proteins RNF28/MURF1 and RNF29/MURF2. In vitro studies demonstrated that this protein, RNF28, and RNF29 form heterodimers, which may be important for the regulation of titin kinase and microtubule-dependent signal pathways in striated muscles. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq].


References

Centner, T., et al. J. Mol. Biol. 306(4):717-726(2001) Spencer, J.A., et al. J. Cell Biol. 150(4):771-784(2000)

Images



TRIM54 Antibody (N-term) (Cat. #AP13292a) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the TRIM54 antibody detected the TRIM54 protein (arrow).



TRIM54 Antibody (N-term) (Cat. #AP13292a)immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TRIM54 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.