

Thioredoxin (TRX) Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP1337a

Product Information

Application	WB, E
Primary Accession	P10599
Other Accession	Q5T937
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB5217
Calculated MW	11737
Antigen Region	1-30

Additional Information

Gene ID	7295
Other Names	Thioredoxin, Trx, ATL-derived factor, ADF, Surface-associated sulphhydryl protein, SASP, TXN, TRDX, TRX, TRX1
Target/Specificity	This Thioredoxin (TRX) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human Thioredoxin (TRX).
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Thioredoxin (TRX) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TXN
Synonyms	TRDX, TRX, TRX1
Function	Participates in various redox reactions through the reversible oxidation of

its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions (PubMed:[17182577](#), PubMed:[19032234](#), PubMed:[2176490](#)). Plays a role in the reversible S- nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide. Nitrosylates the active site Cys of CASP3 in response to nitric oxide (NO), and thereby inhibits caspase-3 activity (PubMed:[16408020](#), PubMed:[17606900](#)). Induces the FOS/JUN AP-1 DNA-binding activity in ionizing radiation (IR) cells through its oxidation/reduction status and stimulates AP-1 transcriptional activity (PubMed:[11118054](#), PubMed:[9108029](#)).

Cellular Location

Nucleus. Cytoplasm. Secreted Note=Translocates from the cytoplasm into the nucleus after phorbol 12- myristate 13-acetate induction (PMA) (PubMed:[9108029](#)). Predominantly in the cytoplasm in non irradiated cells (PubMed:[11118054](#)). Radiation induces translocation of TRX from the cytoplasm to the nucleus (PubMed:[11118054](#)). Secreted by a leaderless secretory pathway (PubMed:[1332947](#)).


Background

Thioredoxins (Trx) are small, multi-functional proteins with oxidoreductase activity and are ubiquitous in essentially all living cells. Trx contains a redox active disulfide/dithiol group within the conserved Cys-Gly-Pro-Cys active site. The two cysteine residues in the conserved active centers can be oxidized to form intramolecular disulfide bonds. Reduction of the active site disulfide in oxidized Trx is catalyzed by Trx reductase with NADPH as the electron donor. The reduced Trx is a hydrogen donor for ribonucleotide reductase, the essential enzyme for DNA synthesis, and a potent general protein disulfide reductase with numerous functions in growth and redox regulations. Specific protein disulfide targets for reduction by Trx include protein disulfide isomerase(PDI) and a number of transcription factors such as p53, NF-kB and AP-1. Trx is also capable of removing H₂O₂, particularly when it is coupled with either methionine sulfoxide reductase or several isoforms of peroxiredoxins.

References


Cell. Signal. 17 (8), 985-996 (2005)
 Redox Rep. 29 (3), 281-286 (2005)
 Blood 105 (4), 1598-1605 (2005)
 Oncogene 23 (55), 8868-8875 (2004)
 J. Biol. Chem. 279 (29), 30369-30374 (2004)

Images



Anti-TrX Antibody (M1) at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 12 kDa
 Blocking/Dilution buffer: 5% NFDM/TBST.

Western blot analysis of anti-TrX Pab (AP1337a) in HL-60 cell line lysate. TrX(arrow) was detected using the purified Pab.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.