

CTGF Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13399C

Product Information

Application	WB, IHC-P, E
Primary Accession	P29279
Other Accession	Q9R1E9 , O19113 , P29268 , NP_001892.1
Reactivity	Human, Rat, Mouse
Predicted	Mouse, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB33090
Calculated MW	38091
Antigen Region	166-193

Additional Information

Gene ID	1490
Other Names	Connective tissue growth factor, CCN family member 2, Hypertrophic chondrocyte-specific protein 24, Insulin-like growth factor-binding protein 8, IBP-8, IGF-binding protein 8, IGFBP-8, CTGF, CCN2, HCS24, IGFBP8
Target/Specificity	This CTGF antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 166-193 amino acids from the Central region of human CTGF.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CTGF Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CCN2 (HGNC:2500)
Function	Major connective tissue chemoattractant secreted by vascular endothelial

cells. Promotes proliferation and differentiation of chondrocytes. Is involved in the stimulation of osteoblast differentiation and has a critical role in osteogenesis (PubMed:[39414788](#)). Mediates heparin- and divalent cation-dependent cell adhesion in many cell types including fibroblasts, myofibroblasts, endothelial and epithelial cells. Enhances fibroblast growth factor- induced DNA synthesis.

Cellular Location

Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:P29268}. Secreted

Tissue Location

Expressed in bone marrow and thymic cells. Also expressed one of two Wilms tumors tested.


Background

The protein encoded by this gene is a mitogen that is secreted by vascular endothelial cells. The encoded protein plays a role in chondrocyte proliferation and differentiation, cell adhesion in many cell types, and is related to platelet-derived growth factor. Certain polymorphisms in this gene have been linked with a higher incidence of systemic sclerosis. [provided by RefSeq].


References

Behrens, M.E., et al. *Oncogene* 29(42):5667-5677(2010)
Cunningham, J.L., et al. *Eur. J. Endocrinol.* 163(4):691-697(2010)
Ito, Y., et al. *Am. J. Physiol. Renal Physiol.* 299 (3), F545-F558 (2010) :
Adler, S.G., et al. *Clin J Am Soc Nephrol* 5(8):1420-1428(2010)
Johnatty, S.E., et al. *PLoS Genet.* 6 (7), E1001016 (2010) :

Images



All lanes : Anti-CTGF Antibody (Center) at 1:1000 dilution
Lane 1: HUVEC whole cell lysate Lane 2: mouse heart lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 38kDa
Blocking/Dilution buffer : 5% NFDM/TBST.



CTGF Antibody (Center) (Cat. #AP13399c) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CTGF Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.