

TRIM39 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13466c

Product Information

Application	WB, IHC-P, E
Primary Accession	Q9HCM9
Other Accession	NP_742013.1 , NP_067076.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB33252
Calculated MW	59690
Antigen Region	248-277

Additional Information

Gene ID	56658
Other Names	E3 ubiquitin-protein ligase TRIM39, 632-, RING finger protein 23, Testis-abundant finger protein, Tripartite motif-containing protein 39, TRIM39, RNF23, TFP
Target/Specificity	This TRIM39 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 248-277 amino acids from the Central region of human TRIM39.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TRIM39 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TRIM39
Synonyms	RNF23, TFP

Function	[Isoform 1]: E3 ubiquitin-protein ligase (PubMed: 22529100). May facilitate apoptosis by inhibiting APC/C-Cdh1-mediated poly-ubiquitination and subsequent proteasome-mediated degradation of the pro-apoptotic protein MOAP1 (PubMed: 19100260 , PubMed: 22529100). Regulates the G1/S transition of the cell cycle and DNA damage-induced G2 arrest by stabilizing CDKN1A/p21 (PubMed: 23213251). Positively regulates CDKN1A/p21 stability by competing with DTL for CDKN1A/p21 binding, therefore disrupting DCX(DTL) E3 ubiquitin ligase complex- mediated CDKN1A/p21 ubiquitination and degradation (PubMed: 23213251).
Cellular Location	[Isoform 1]: Cytoplasm, cytosol. Mitochondrion. Nucleus Note=Found predominantly in the cytosol. Partial shift from the cytosol to the mitochondria when colocalized with MOAP1. Colocalizes with CDKN1A in the nucleus.
Tissue Location	Ubiquitous; highly expressed in brain, heart, kidney, liver, skeletal muscle, spleen and testis


Background

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The function of this protein has not been identified. This gene lies within the major histocompatibility complex class I region on chromosome 6. Alternate splicing results in two transcript variants encoding different isoforms. [provided by RefSeq].

References


Kurata, R., et al. Biochem. Biophys. Res. Commun. 401(4):533-537(2010)
 Barcellos, L.F., et al. PLoS Genet. 5 (10), E1000696 (2009) :
 Lee, S.S., et al. Exp. Cell Res. 315(7):1313-1325(2009)
 Roberts, J.D. Jr., et al. Am. J. Physiol. Lung Cell Mol. Physiol. 293 (4), L903-L912 (2007) :
 Wu, C., et al. Proteomics 7(11):1775-1785(2007)

Images



TRIM39 Antibody (Center) (Cat. #AP13466c) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the TRIM39 antibody detected the TRIM39 protein (arrow).

TRIM39 Antibody (Center) (Cat. #AP13466c) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TRIM39 Antibody (Center) for immunohistochemistry. Clinical relevance has not been



evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.