10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ZBTB11 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13526b

Product Information

Application	WB, E
Primary Accession	O95625
Other Accession	NP_055230.2
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB33568
Calculated MW	119384
Antigen Region	736-765

Additional Information

Gene ID	27107
Other Names	Zinc finger and BTB domain-containing protein 11, ZBTB11
Target/Specificity	This ZBTB11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 736-765 amino acids from the C-terminal region of human ZBTB11.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ZBTB11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ZBTB11 {ECO:0000303|PubMed:28397838, ECO:0000312|HGNC:HGNC:16740}
Function	May be involved in transcriptional regulation.

Cellular Location Nucleus, nucleolus.


Background

ZBTB11 may be involved in transcriptional regulation.

References

Colland, F., et al. Genome Res. 14(7):1324-1332(2004)

Images



All lanes : Anti-ZBTB11 Antibody (C-term) at 1:1000 dilution Lane 1: A431 whole cell lysate Lane 2: Hela whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 119 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.