10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TTC13 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13674a

Product Information

Application	WB, IHC-P, E
Primary Accession	Q8NBP0
Other Accession	NP_078801.3, NP_001116307.2
Reactivity	Human
Predicted	Mouse, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB33472
Calculated MW	96813
Antigen Region	195-224

Additional Information

Gene ID	79573
Other Names	Tetratricopeptide repeat protein 13, TPR repeat protein 13, TTC13
Target/Specificity	This TTC13 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 195-224 amino acids from the N-terminal region of human TTC13.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TTC13 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TTC13


Background

The function of this protein remains unknown.


References

Kimura, K., et al. Genome Res. 16(1):55-65(2006)

Images



TTC13 Antibody (N-term) (Cat. #AP13674a) western blot analysis in U251 cell line lysates (35ug/lane).This demonstrates the TTC13 antibody detected the TTC13 protein (arrow).



TTC13 Antibody (N-term) (Cat. #AP13674a)immunohistochemistry analysis in formalin fixed and paraffin embedded human cerebellum tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of TTC13 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Citations

- TTC13 expression and STAT3 activation may form a positive feedback loop to promote ccRCC progression

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.