10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# CSK Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP13748a

## Product Information

| | |
|---|---|
| **Application** | WB, E |
| **Primary Accession** | P41240 |
| **Other Accession** | P32577, P41241, P41239, Q0VBZ0, NP_001120662.1, NP_004374.1 |
| **Reactivity** | Human |
| **Predicted** | Bovine, Chicken, Mouse, Rat, Canine |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB33742 |
| **Calculated MW** | 50704 |
| **Antigen Region** | 33-61 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 1445 |
| **Other Names** | Tyrosine-protein kinase CSK, C-Src kinase, Protein-tyrosine kinase CYL, CSK |
| **Target/Specificity** | This CSK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 33-61 amino acids from the N-terminal region of human CSK. |
| **Dilution** | WB~~1:1000 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | CSK Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | CSK |
| **Function** | Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C- terminal tails of Src-family |

| | |
|---|---|
| | kinases (SFKs) including LCK, SRC, HCK, FYN, LYN, CSK or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T-cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK. May act as a negative regulator of EGFR and STAT3 signaling pathways (PubMed:26918609). |
| **Cellular Location** | Cytoplasm {ECO:0000250\|UniProtKB:P41241}. Cell membrane {ECO:0000250\|UniProtKB:P41241}. Note=Mainly cytoplasmic, also present in lipid rafts. {ECO:0000250\|UniProtKB:P41241} |
| **Tissue Location** | Expressed in lung and macrophages. |


## Background

Specifically phosphorylates 'Tyr-504' on LCK, which acts as a negative regulatory site. Can also act on the LYN and FYN kinases.

## References


Niu, W., et al. Clin. Chim. Acta 411 (19-20), 1491-1495 (2010) :
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)
Mikkola, E.T., et al. J. Mol. Biol. 399(4):618-627(2010)
Takeuchi, F., et al. Circulation 121(21):2302-2309(2010)


## Images




CSK Antibody (N-term) (Cat. #AP13748a) western blot analysis in Hela cell line lysates (35ug/lane).This demonstrates the CSK antibody detected the CSK protein (arrow).

## Citations

- Overexpression of Csk-binding protein/phosphoprotein associated with glycosphingolipid-enriched microdomains induces cluster of differentiation 59-mediated apoptosis in Jurkat cells.


Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.