

CBX5 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13779c

Product Information

Application	WB, E
Primary Accession	P45973
Other Accession	Q61686 , NP_036249.1 , NP_001120794.1 , NP_001120793.1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB33905
Calculated MW	22225
Antigen Region	88-117

Additional Information

Gene ID	23468
Other Names	Chromobox protein homolog 5, Antigen p25, Heterochromatin protein 1 homolog alpha, HP1 alpha, CBX5, HP1A
Target/Specificity	This CBX5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 88-117 amino acids from the Central region of human CBX5.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CBX5 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CBX5
Synonyms	HP1A

Function Component of heterochromatin that recognizes and binds histone H3 tails methylated at 'Lys-9' (H3K9me), leading to epigenetic repression. In contrast, it is excluded from chromatin when 'Tyr-41' of histone H3 is phosphorylated (H3Y41ph) (PubMed:[19783980](#)). May contribute to the association of heterochromatin with the inner nuclear membrane by interactions with the lamin-B receptor (LBR) (PubMed:[19783980](#)). Involved in the formation of kinetochore through interaction with the MIS12 complex subunit NSL1 (PubMed:[19783980](#), PubMed:[20231385](#)). Required for the formation of the inner centromere (PubMed:[20231385](#)).

Cellular Location Nucleus. Chromosome. Chromosome, centromere. Note=Colocalizes with HNRNPU in the nucleus (PubMed:19617346). Component of centromeric and pericentromeric heterochromatin. Associates with chromosomes during mitosis. Associates specifically with chromatin during metaphase and anaphase (PubMed:19617346). Localizes to sites of DNA damage (PubMed:28977666)


Background

This gene encodes a highly conserved nonhistone protein, which is a member of the heterochromatin protein family. The protein is enriched in the heterochromatin and associated with centromeres. The protein has a single N-terminal chromodomain which can bind to histone proteins via methylated lysine residues, and a C-terminal chromo shadow-domain (CSD) which is responsible for the homodimerization and interaction with a number of chromatin-associated nonhistone proteins. The encoded product is involved in the formation of functional kinetochore through interaction with essential kinetochore proteins. The gene has a pseudogene located on chromosome 3. Multiple alternatively spliced variants, encoding the same protein, have been identified.

References

Nozawa, R.S., et al. Nat. Cell Biol. 12(7):719-727(2010)
Zeng, W., et al. Epigenetics 5(4):287-292(2010)
Emelyanov, A.V., et al. J. Biol. Chem. 285(20):15027-15037(2010)
Kiyomitsu, T., et al. J. Cell Biol. 188(6):791-807(2010)
Chaturvedi, P., et al. PLoS ONE 5 (5), E10620 (2010) :

Images



CBX5 Antibody (Center) (Cat. #AP13779c) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the CBX5 antibody detected the CBX5 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.