

# DHRS7C Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13788c

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">A6NNS2</a>
<b>Other Accession</b>	<a href="#">Q8CHS7</a> , <a href="#">Q1RMJ5</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Bovine, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB33974
<b>Calculated MW</b>	34878
<b>Antigen Region</b>	116-145

## Additional Information

---

<b>Gene ID</b>	201140
<b>Other Names</b>	Dehydrogenase/reductase SDR family member 7C, 11--, DHRS7C
<b>Target/Specificity</b>	This DHRS7C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 116-145 amino acids from the Central region of human DHRS7C.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	DHRS7C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	DHRS7C ( <a href="#">HGNC:32423</a> )
<b>Function</b>	NADH-dependent oxidoreductase which catalyzes the oxidation of all-trans-retinol to all-trans-retinal. Plays a role in the regulation of cardiac and skeletal muscle metabolic functions. Maintains Ca(2+) intracellular

homeostasis by repressing  $\text{Ca}^{2+}$  release from the sarcoplasmic reticulum (SR) in myotubes, possibly through local alternations in NAD/NADH or retinol/retinal. Also plays a role in  $\text{Ca}^{2+}$  homeostasis by controlling  $\text{Ca}^{2+}$  overload in the cytosol and the SR in myotubes. Involved in glucose uptake into skeletal muscles and muscle performance by activating PI3K and mTORC2-mediated AKT1 phosphorylation signaling pathways, possibly through the action of its downstream catalytic product all-trans-retinoic acid.


### Cellular Location

Sarcoplasmic reticulum membrane {ECO:0000250 | UniProtKB:Q8CHS7}. Note=The N-terminus region encompasses a short hydrophobic sequence bound to the sarcoplasmic reticulum membrane, whereas the C-terminus catalytic domain faces the myoplasm In skeletal muscle, enriched in the longitudinal sarcoplasmic reticulum. {ECO:0000250 | UniProtKB:Q8CHS7}


## Background

DHRS7C is a putative oxidoreductase (Potential).

## Images



DHRS7C Antibody (Center) (Cat. #AP13788c) western blot analysis in human placenta tissue lysates (35ug/lane). This demonstrates the DHRS7C antibody detected the DHRS7C protein (arrow).



DHRS7C Antibody (Center) (AP13788c) immunohistochemistry analysis in formalin fixed and paraffin embedded human heart tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of DHRS7C Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.