

WNT8B Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13948b

Product Information

Application	WB, E
Primary Accession	Q93098
Other Accession	NP_003384.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB22565
Calculated MW	38721
Antigen Region	323-351

Additional Information

Gene ID	7479
Other Names	Protein Wnt-8b, WNT8B
Target/Specificity	This WNT8B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 323-351 amino acids from the C-terminal region of human WNT8B.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WNT8B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	WNT8B
Function	Ligand for members of the frizzled family of seven transmembrane receptors. May play an important role in the development and differentiation of certain forebrain structures, notably the hippocampus.

Cellular Location	Secreted, extracellular space, extracellular matrix
Tissue Location	Expression is restricted to the brain, and more specifically to the forebrain.


Background

The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It encodes a protein which shows 95%, 86% and 71% amino acid identity to the mouse, zebrafish and *Xenopus* Wnt8B proteins, respectively. The expression patterns of the human and mouse genes appear identical and are restricted to the developing brain. The chromosomal location of this gene to 10q24 suggests it as a candidate gene for partial epilepsy.

References

Katoh, M., et al. *Int. J. Oncol.* 30(5):1273-1277(2007)
 Grupe, A., et al. *Am. J. Hum. Genet.* 78(1):78-88(2006)
 Deloukas, P., et al. *Nature* 429(6990):375-381(2004)
 Saitoh, T., et al. *Int. J. Oncol.* 20(5):999-1003(2002)
 Saitoh, T., et al. *Int. J. Oncol.* 20(2):343-348(2002)

Images



Anti-WNT8B Antibody (C-term) at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.