

ITGB5 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14000a

Product Information

Application	WB, IHC-P, E
Primary Accession	P18084
Other Accession	O70309 , P80747 , NP_002204.2
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB33832
Calculated MW	88054
Antigen Region	216-244

Additional Information

Gene ID	3693
Other Names	Integrin beta-5, ITGB5
Target/Specificity	This ITGB5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 216-244 amino acids from the N-terminal region of human ITGB5.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ITGB5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ITGB5
Function	Integrin alpha-V/beta-5 (ITGAV:ITGB5) is a receptor for fibronectin. It recognizes the sequence R-G-D in its ligand.


Background

Integrin alpha-V/beta-5 is a receptor for fibronectin. It recognizes the sequence R-G-D in its ligand.


References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Li, Z., et al. J. Biol. Chem. 285(31):23699-23710(2010)
Leifheit-Nestler, M., et al. Arterioscler. Thromb. Vasc. Biol. 30(7):1398-1406(2010)
Lane, D., et al. Oncogene 29(24):3519-3531(2010)
Lyle, C., et al. Virol. J. 7, 148 (2010) :

Images



ITGB5 Antibody (N-term) (Cat. #AP14000a) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the ITGB5 antibody detected the ITGB5 protein (arrow).



ITGB5 Antibody (N-term) (Cat. #AP14000a) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of ITGB5 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Citations

- [Cisplatin suppresses the growth and proliferation of breast and cervical cancer cell lines by inhibiting integrin \$\beta\$ 5-mediated glycolysis.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.