

# NOX4 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14025A

## Product Information

---

Application	IHC-P, WB, E
Primary Accession	<a href="#">Q9NPH5</a>
Other Accession	<a href="#">NP_001137308.1</a> , <a href="#">NP_058627.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB34043
Calculated MW	66932
Antigen Region	124-153

## Additional Information

---

Gene ID	50507
Other Names	NADPH oxidase 4, 163-, Kidney oxidase-1, KOX-1, Kidney superoxide-producing NADPH oxidase, Renal NAD(P)H-oxidase, NOX4, RENOX
Target/Specificity	This NOX4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 124-153 amino acids from the N-terminal region of human NOX4.
Dilution	IHC-P~~1:100~500 WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NOX4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	NOX4
Synonyms	RENOX
Function	NADPH oxidase that catalyzes predominantly the reduction of oxygen to

H2O2 (PubMed:[14966267](#), PubMed:[15356101](#), PubMed:[15927447](#), PubMed:[21343298](#), PubMed:[25062272](#)). Can also catalyze to a smaller extent, the reduction of oxygen to superoxide (PubMed:[10869423](#), PubMed:[11032835](#), PubMed:[15155719](#), PubMed:[15572675](#), PubMed:[15927447](#), PubMed:[16019190](#), PubMed:[16179589](#), PubMed:[16230378](#), PubMed:[16324151](#), PubMed:[25062272](#)). May function as an oxygen sensor regulating the KCNK3/TASK-1 potassium channel and HIF1A activity (PubMed:[16019190](#)). May regulate insulin signaling cascade (PubMed:[14966267](#)). May play a role in apoptosis, bone resorption and lipopolysaccharide-mediated activation of NFkB (PubMed:[15356101](#), PubMed:[15572675](#)). May produce superoxide in the nucleus and play a role in regulating gene expression upon cell stimulation (PubMed:[16324151](#)). Promotes ferroptosis, reactive oxygen species production and reduced glutathione (GSH) levels by activating NLRP3 inflammasome activation and cytokine release (PubMed:[39909992](#)).

#### Cellular Location

Cytoplasm. Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Cell junction, focal adhesion {ECO:0000250|UniProtKB:Q924V1}. Nucleus [Isoform 3]: Cytoplasm. Cytoplasm, perinuclear region [Isoform 6]: Cytoplasm. Cytoplasm, perinuclear region

#### Tissue Location

Expressed by distal tubular cells in kidney cortex and in endothelial cells (at protein level). Widely expressed. Strongly expressed in kidney and to a lower extent in heart, adipocytes, hepatoma, endothelial cells, skeletal muscle, brain, several brain tumor cell lines and airway epithelial cells

## Background

---

This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localized to non-phagocytic cells where it acts as an oxygen sensor and catalyzes the reduction of molecular oxygen to various reactive oxygen species (ROS). The ROS generated by this protein have been implicated in numerous biological functions including signal transduction, cell differentiation and tumor cell growth. A pseudogene has been identified on the other arm of chromosome 11. Alternative splicing results in multiple transcript variants.

## References



---

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)  
 Salmeen, A., et al. Oncogene 29(31):4473-4484(2010)  
 Wu, R.F., et al. Mol. Cell. Biol. 30(14):3553-3568(2010)  
 Diebold, I., et al. Mol. Biol. Cell 21(12):2087-2096(2010)  
 Manea, A., et al. Biochem. Biophys. Res. Commun. 396(4):901-907(2010)

## Images

---

NOX4 Antibody (N-term) (Cat. #AP14025a) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the NOX4 antibody detected the NOX4 protein (arrow).



NOX4 Antibody (N-term)  
(AP14025a) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of NOX4 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.