

TICAM2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14060a

Product Information

Application	WB, E
Primary Accession	Q86XR7
Other Accession	NP_001157941.1 , NP_067681.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB34146
Calculated MW	26916
Antigen Region	12-41

Additional Information

Gene ID	100302736;353376
Other Names	TIR domain-containing adapter molecule 2, TICAM-2, Putative NF-kappa-B-activating protein 502, TRIF-related adapter molecule, Toll-like receptor adaptor protein 3, Toll/interleukin-1 receptor domain-containing protein, MyD88-4, TICAM2, TIRAP3, TIRP, TRAM
Target/Specificity	This TICAM2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 12-41 amino acids from the N-terminal region of human TICAM2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TICAM2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TICAM2
Synonyms	TIRAP3, TIRP, TRAM

Function	Functions as a sorting adapter in different signaling pathways to facilitate downstream signaling leading to type I interferon induction (PubMed: 16603631 , PubMed: 16757566 , PubMed: 25385819 , PubMed: 25825441). In TLR4 signaling, physically bridges TLR4 and TICAM1 and functionally transmits signal to TICAM1 in early endosomes after endocytosis of TLR4. In TLR2 signaling, physically bridges TLR2 and MYD88 and is required for the TLR2- dependent movement of MYD88 to endosomes following ligand engagement (PubMed: 25385819). Involved in IL-18 signaling and is proposed to function as a sorting adapter for MYD88 in IL-18 signaling during adaptive immune response (PubMed: 22685567). Forms a complex with RAB11FIP2 that is recruited to the phagosomes to promote the activation of the actin-regulatory GTPases RAC1 and CDC42 and subsequent phagocytosis of Gram-negative bacteria (PubMed: 30883606).
Cellular Location	[Isoform 1]: Cytoplasm. Golgi apparatus. Cell membrane. Endoplasmic reticulum. Early endosome membrane. Late endosome membrane. Cell projection, phagocytic cup. Note=Localized to the plasma membrane as a result of myristoylation. Phosphorylation on Ser-16 leads to its depletion from the membrane. Upon LPS stimulation colocalizes with isoform 2 in late endosomes
Tissue Location	Expressed in spleen, prostate, testis, uterus, small intestine, colon, peripheral blood leukocytes, heart, placenta, lung, liver, skeletal muscle, and pancreas Isoform 2 is ubiquitously expressed (at lower levels than isoform 1)


Background

TIRP is a Toll/interleukin-1 receptor (IL1R; MIM 147810) (TIR) domain-containing adaptor protein involved in Toll receptor signaling (see TLR4; MIM 603030).

References

Lysakova-Devine, T., et al. J. Immunol. 185(7):4261-4271(2010)
 Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
 Palsson-McDermott, E.M., et al. Nat. Immunol. 10(6):579-586(2009)
 Hawn, T.R., et al. PLoS ONE 4 (6), E5990 (2009) :
 Nakajima, T., et al. Immunogenetics 60(12):727-735(2008)

Images



TICAM2 Antibody (N-term) (Cat. #AP14060a) western blot analysis in human placenta tissue lysates (35ug/lane). This demonstrates the TICAM2 antibody detected the TICAM2 protein (arrow).