

IGF1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14099b

Product Information

Application WB, IF, FC, IHC-P, E

Primary Accession P05019

Other Accession <u>P01343</u>, <u>NP 000609.1</u>

Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB33711
Calculated MW 21841
Antigen Region 115-144

Additional Information

Gene ID 3479

Other Names IGF1; IBP1; Insulin-like growth factor I; Mechano growth factor;

Somatomedin-C

Target/Specificity This IGF1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 115-144 amino acids from the

C-terminal region of human IGF1.

Dilution WB~~1:1000 IF~~1:10~50 FC~~1:10~50 IHC-P~~1:100~500 E~~Use at an assay

dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This

antibody is purified through a protein A column, followed by peptide affinity

purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions IGF1 Antibody (C-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name IGF1 (<u>HGNC:5464</u>)

Function The insulin-like growth factors, isolated from plasma, are structurally and

functionally related to insulin but have a much higher growth-promoting

activity. May be a physiological regulator of [1-14C]- 2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblasts. Stimulates glucose transport in bone-derived osteoblastic (PyMS) cells and is effective at much lower concentrations than insulin, not only regarding glycogen and DNA synthesis but also with regard to enhancing glucose uptake. May play a role in synapse maturation (PubMed:21076856, PubMed:24132240). Ca(2+)-dependent exocytosis of IGF1 is required for sensory perception of smell in the olfactory bulb (By similarity). Acts as a ligand for IGF1R. Binds to the alpha subunit of IGF1R, leading to the activation of the intrinsic tyrosine kinase activity which autophosphorylates tyrosine residues in the beta subunit thus initiating a cascade of down-stream signaling events leading to activation of the PI3K-AKT/PKB and the Ras-MAPK pathways. Binds to integrins ITGAV:ITGB3 and ITGA6:ITGB4. Its binding to integrins and subsequent ternary complex formation with integrins and IGFR1 are essential for IGF1 signaling. Induces the phosphorylation and activation of IGFR1, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (PubMed: 19578119, PubMed: 22351760, PubMed: 23243309, PubMed:23696648). As part of the MAPK/ERK signaling pathway, acts as a negative regulator of apoptosis in cardiomyocytes via promotion of STUB1/CHIP-mediated ubiquitination and degradation of ICER-type isoforms of CREM (By similarity).

Cellular Location

Secreted {ECO:0000250 | UniProtKB:P05017}.


Background

The protein encoded by this gene is similar to insulin in function and structure and is a member of a family of proteins involved in mediating growth and development. The encoded protein is processed from a precursor, bound by a specific receptor, and secreted. Defects in this gene are a cause of insulin-like growth factor I deficiency. Several transcript variants encoding different isoforms have been found for this gene.

References


Li, M., et al. J. Biol. Chem. 285(40):30480-30488(2010) Canzian, F., et al. Hum. Mol. Genet. 19(19):3873-3884(2010) Romero, R., et al. Am. J. Obstet. Gynecol. 203 (4), 361 (2010): Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Szczesny, G., et al. Arch Orthop Trauma Surg (2010) In press:


Images




All lanes: Anti- IGF1 Antibody (C-term) at 1:1000 dilution + SiHa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 22 KDa Blocking/Dilution buffer: 5% NFDM/TBST.


Confocal immunofluorescent analysis of IGF1 Antibody (C-term)(Cat#AP14099b) with A549 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green).DAPI







IGF1 Antibody (C-term) (Cat. #AP14099b) flow cytometric analysis of A549 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



IGF1 Antibody (C-term) (Cat. #AP14099b)immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of IGF1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Citations

• The Role of MiR-5094 as a Proliferation Suppressor during Cellular Radiation Response via Downregulating STAT5b

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.