

TCEB3 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14198c

Product Information

Application WB, E **Primary Accession** Q14241 Other Accession NP 003189.2 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB33959 **Calculated MW** 87230 253-282 **Antigen Region**

Additional Information

Gene ID 6924

Other Names Transcription elongation factor B polypeptide 3, Elongin 110 kDa subunit,

Elongin-A, EloA, RNA polymerase II transcription factor SIII subunit A1, SIII

p110, TCEB3

Target/Specificity This TCEB3 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 253-282 amino acids from the Central

region of human TCEB3.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions TCEB3 Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name ELOA (HGNC:11620)

Synonyms TCEB3

Function SIII, also known as elongin, is a general transcription elongation factor that

increases the RNA polymerase II transcription elongation past

template-encoded arresting sites. Subunit A is transcriptionally active and its transcription activity is strongly enhanced by binding to the dimeric complex

of the SIII regulatory subunits B and C (elongin BC complex).

Cellular Location Nucleus. Note=Localizes to sites of DNA damage.


Background

This gene encodes the protein elongin A, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation.

References

Landa, I., et al. PLoS Genet. 5 (9), E1000637 (2009):
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Woo, J.S., et al. EMBO J. 25(6):1353-1363(2006)
Andersen, J.S., et al. Nature 433(7021):77-83(2005)
Beausoleil, S.A., et al. Proc. Natl. Acad. Sci. U.S.A. 101(33):12130-12135(2004)

Images



TCEB3 Antibody (Center) (Cat. #AP14198c) western blot analysis in Hela cell line lysates (35ug/lane). This demonstrates the TCEB3 antibody detected the TCEB3 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.