

TCEB3 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14198c

Product Information

Application	WB, E
Primary Accession	Q14241
Other Accession	NP_003189.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB33959
Calculated MW	87230
Antigen Region	253-282

Additional Information

Gene ID	6924
Other Names	Transcription elongation factor B polypeptide 3, Elongin 110 kDa subunit, Elongin-A, EloA, RNA polymerase II transcription factor SIII subunit A1, SIII p110, TCEB3
Target/Specificity	This TCEB3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 253-282 amino acids from the Central region of human TCEB3.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TCEB3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ELOA (HGNC:11620)
Synonyms	TCEB3

Function	SIII, also known as elongin, is a general transcription elongation factor that increases the RNA polymerase II transcription elongation past template-encoded arresting sites. Subunit A is transcriptionally active and its transcription activity is strongly enhanced by binding to the dimeric complex of the SIII regulatory subunits B and C (elongin BC complex).
Cellular Location	Nucleus. Note=Localizes to sites of DNA damage.


Background

This gene encodes the protein elongin A, which is a subunit of the transcription factor B (SIII) complex. The SIII complex is composed of elongins A/A2, B and C. It activates elongation by RNA polymerase II by suppressing transient pausing of the polymerase at many sites within transcription units. Elongin A functions as the transcriptionally active component of the SIII complex, whereas elongins B and C are regulatory subunits. Elongin A2 is specifically expressed in the testis, and capable of forming a stable complex with elongins B and C. The von Hippel-Lindau tumor suppressor protein binds to elongins B and C, and thereby inhibits transcription elongation.

References

Landa, I., et al. PLoS Genet. 5 (9), E1000637 (2009) :
 Olsen, J.V., et al. Cell 127(3):635-648(2006)
 Woo, J.S., et al. EMBO J. 25(6):1353-1363(2006)
 Andersen, J.S., et al. Nature 433(7021):77-83(2005)
 Beausoleil, S.A., et al. Proc. Natl. Acad. Sci. U.S.A. 101(33):12130-12135(2004)

Images



TCEB3 Antibody (Center) (Cat. #AP14198c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the TCEB3 antibody detected the TCEB3 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.