

CYTH1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14202a

Product Information

Application	WB, E
Primary Accession	Q15438
Other Accession	P97694 , Q9QX11 , NP_004753.1 , NP_059430.2
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB34095
Calculated MW	46413
Antigen Region	37-65

Additional Information

Gene ID	9267
Other Names	Cytohesin-1, PH, SEC7 and coiled-coil domain-containing protein 1, SEC7 homolog B2-1, CYTH1, D17S811E, PSCD1
Target/Specificity	This CYTH1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 37-65 amino acids from the N-terminal region of human CYTH1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CYTH1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CYTH1 (HGNC:9501)
Synonyms	D17S811E, PSCD1

Function	Promotes guanine-nucleotide exchange on ARF1, ARF5 and ARF6. Promotes the activation of ARF factors through replacement of GDP with GTP. Plays an important role in membrane trafficking, during junctional remodeling and epithelial polarization, through regulation of ARF6 activity.
Cellular Location	Cell membrane; Peripheral membrane protein. Cytoplasm, cytosol {ECO:0000250 UniProtKB:Q9QX11}. Cell junction, tight junction {ECO:0000250 UniProtKB:Q9QX11}. Cell junction, adherens junction {ECO:0000250 UniProtKB:Q9QX11}. Note=Colocalized with TJP1 during epithelial polarization. {ECO:0000250 UniProtKB:Q9QX11}
Tissue Location	Ubiquitous.


Background

The protein encoded by this gene is a member of the PSCD family. Members of this family have identical structural organization that consists of an N-terminal coiled-coil motif, a central Sec7 domain, and a C-terminal pleckstrin homology (PH) domain. The coiled-coil motif is involved in homodimerization, the Sec7 domain contains guanine-nucleotide exchange protein (GEP) activity, and the PH domain interacts with phospholipids and is responsible for association of PSCDs with membranes. Members of this family appear to mediate the regulation of protein sorting and membrane trafficking. This gene is highly expressed in natural killer and peripheral T cells, and regulates the adhesiveness of integrins at the plasma membrane of lymphocytes. The encoded protein is 83% homologous to that of CYTH2.

References

El Azreq, M.A., et al. J. Immunol. 184(2):637-649(2010)
Quast, T., et al. Blood 113(23):5801-5810(2009)
Sendide, K., et al. J. Immunol. 174(7):4210-4219(2005)
Boehm, T., et al. EMBO J. 22(5):1014-1024(2003)
Mansour, M., et al. J. Biol. Chem. 277(35):32302-32309(2002)

Images



CYTH1 Antibody (N-term) (Cat. #AP14202a) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the CYTH1 antibody detected the CYTH1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.