

# NR2F2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14204a

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P24468</a>
<b>Other Accession</b>	<a href="#">O09018</a> , <a href="#">P43135</a> , <a href="#">Q9TTR7</a> , <a href="#">NP_066285.1</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Predicted</b>	Bovine, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	45571
<b>Antigen Region</b>	1-30

## Additional Information

---

<b>Gene ID</b>	7026
<b>Other Names</b>	COUP transcription factor 2, COUP-TF2, Apolipoprotein A-I regulatory protein 1, ARP-1, COUP transcription factor II, COUP-TF II, Nuclear receptor subfamily 2 group F member 2, NR2F2, ARP1, TFCOUP2
<b>Target/Specificity</b>	This NR2F2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human NR2F2.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	NR2F2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	NR2F2
<b>Synonyms</b>	ARP1, TFCOUP2

<b>Function</b>	Ligand-activated transcription factor. Activated by high concentrations of 9-cis-retinoic acid and all-trans-retinoic acid, but not by dexamethasone, cortisol or progesterone (in vitro). Regulation of the apolipoprotein A-I gene transcription. Binds to DNA site A. May be required to establish ovary identity during early gonad development (PubMed: <a href="#">29478779</a> ).
<b>Cellular Location</b>	Nucleus.
<b>Tissue Location</b>	Ubiquitous. Expressed in the stromal cells of developing fetal ovaries (PubMed:29478779)


## Background

This gene encodes a member of the steroid thyroid hormone superfamily of nuclear receptors. The encoded protein is a ligand inducible transcription factor that is involved in the regulation of many different genes. Alternate splicing results in multiple transcript variants.

## References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)  
Feng, T., et al. Hum. Genet. 128(3):269-280(2010)  
Kang, J., et al. Blood 116(1):140-150(2010)  
Eriksson, N., et al. PLoS Genet. 6 (6), E1000993 (2010) :  
Hubert, M.A., et al. PLoS ONE 5 (2), E9417 (2010) :

## Images



NR2F2 Antibody (N-term) (Cat. #AP14204a) western blot analysis in HepG2,mouse NIH/3T3 cell line lysates (35ug/lane).This demonstrates the NR2F2 antibody detected the NR2F2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.