

FZD5 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14210b

Product Information

Application	WB, E
Primary Accession	Q13467
Other Accession	NP_003459.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB34120
Calculated MW	64507
Antigen Region	472-500

Additional Information

Gene ID	7855
Other Names	Frizzled-5, Fz-5, hFz5, FzE5, FZD5, C2orf31
Target/Specificity	This FZD5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 472-500 amino acids from the C-terminal region of human FZD5.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	FZD5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FZD5
Synonyms	C2orf31
Function	Receptor for Wnt proteins (PubMed: 10097073 , PubMed: 20530549 , PubMed: 26908622 , PubMed: 9054360). Functions in the canonical Wnt/beta-

catenin signaling pathway. In vitro activates WNT2, WNT10B, WNT5A, but not WNT2B or WNT4 signaling (By similarity). In neurons, activation by WNT7A promotes formation of synapses (PubMed:[20530549](#)). May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues (Probable). Plays a role in yolk sac angiogenesis and in placental vascularization (By similarity). Plays a role in ocular development (PubMed:[26908622](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q8CHL0}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q9EQD0}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q9EQD0}. Synapse {ECO:0000250|UniProtKB:Q8CHL0}. Perikaryon {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, axon {ECO:0000250|UniProtKB:Q8CHL0}. Note=Localized at the plasma membrane and also found at the Golgi apparatus. {ECO:0000250|UniProtKB:Q9EQD0}


Background

Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD5 protein is believed to be the receptor for the Wnt5A ligand.

References

Moller, M., et al. BMC Infect. Dis. 10, 154 (2010) :
Yerges, L.M., et al. J. Bone Miner. Res. 24(12):2039-2049(2009)
Terabayashi, T., et al. J. Biol. Chem. 284(39):26716-26724(2009)
Kim, J.G., et al. J. Korean Med. Sci. 24(3):443-447(2009)
Lin, Z., et al. J. Biol. Chem. 283(48):33053-33058(2008)

Images



FZD5 Antibody (C-term) (Cat. #AP14210b) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the FZD5 antibody detected the FZD5 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.