

# Mouse Prkca Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14277a

## Product Information

---

<b>Application</b>	WB, FC, IHC-P, E
<b>Primary Accession</b>	<a href="#">P20444</a>
<b>Other Accession</b>	<a href="#">P05696</a> , <a href="#">NP_035231.2</a>
<b>Reactivity</b>	Human, Mouse
<b>Predicted</b>	Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB34864
<b>Calculated MW</b>	76852
<b>Antigen Region</b>	1-30

## Additional Information

---

<b>Gene ID</b>	18750
<b>Other Names</b>	Protein kinase C alpha type, PKC-A, PKC-alpha, Prkca, Pkca
<b>Target/Specificity</b>	This Mouse Prkca antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of mouse Prkca.
<b>Dilution</b>	WB~~1:1000 FC~~1:10~50 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	Mouse Prkca Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	Prkca
<b>Synonyms</b>	Pkca

## Function

Calcium-activated, phospholipid- and diacylglycerol (DAG)- dependent serine/threonine-protein kinase that is involved in positive and negative regulation of cell proliferation, apoptosis, differentiation, migration and adhesion, cardiac hypertrophy, angiogenesis, platelet function and inflammation, by directly phosphorylating targets such as RAF1, BCL2, CSPG4, TNNT2/CTNT, or activating signaling cascades involving MAPK1/3 (ERK1/2) and RAP1GAP. Depending on the cell type, is involved in cell proliferation and cell growth arrest by positive and negative regulation of the cell cycle. Can promote cell growth by phosphorylating and activating RAF1, which mediates the activation of the MAPK/ERK signaling cascade, and/or by up-regulating CDKN1A, which facilitates active cyclin-dependent kinase (CDK) complex formation. In cells stimulated by the phorbol ester PMA, can trigger a cell cycle arrest program which is associated with the accumulation of the hyper-phosphorylated growth-suppressive form of RB1 and induction of the CDK inhibitors CDKN1A and CDKN1B. Depending on the cell type, exhibits anti-apoptotic function and protects cells from apoptosis by suppressing the p53/TP53-mediated activation of IGFBP3, or mediates anti-apoptotic action by phosphorylating BCL2. During macrophage differentiation induced by macrophage colony-stimulating factor (CSF1), is translocated to the nucleus and is associated with macrophage development. After wounding, translocates from focal contacts to lamellipodia and participates in the modulation of desmosomal adhesion. Plays a role in cell motility by phosphorylating CSPG4, which induces association of CSPG4 with extensive lamellipodia at the cell periphery and polarization of the cell accompanied by increases in cell motility. During chemokine-induced CD4(+) T cell migration, phosphorylates CDC42-guanine exchange factor DOCK8 resulting in its dissociation from LRCH1 and the activation of GTPase CDC42 (By similarity). Negatively regulates myocardial contractility and positively regulates angiogenesis, platelet aggregation and thrombus formation in arteries. Mediates hypertrophic growth of neonatal cardiomyocytes, in part through a MAPK1/3 (ERK1/2)-dependent signaling pathway, and upon PMA treatment, is required to induce cardiomyocyte hypertrophy up to heart failure and death, by increasing protein synthesis, protein-DNA ratio and cell surface area. Regulates cardiomyocyte function by phosphorylating cardiac troponin T (TNNT2/CTNT), which induces significant reduction in actomyosin ATPase activity, myofilament calcium sensitivity and myocardial contractility. In angiogenesis, is required for full endothelial cell migration, adhesion to vitronectin (VTN), and vascular endothelial growth factor A (VEGFA)-dependent regulation of kinase activation and vascular tube formation. Involved in the stabilization of VEGFA mRNA at post-transcriptional level and mediates VEGFA-induced cell proliferation. In the regulation of calcium-induced platelet aggregation, mediates signals from the CD36/GP4 receptor for granule release, and activates the integrin heterodimer ITGA2B-ITGB3 through the RAP1GAP pathway for adhesion. During response to lipopolysaccharides (LPS), may regulate selective LPS-induced macrophage functions involved in host defense and inflammation. But in some inflammatory responses, may negatively regulate NF-kappa-B-induced genes, through IL1A-dependent induction of NF-kappa-B inhibitor alpha (NFKBIA/IKBA). Upon stimulation with 12-O-tetradecanoylphorbol-13-acetate (TPA), phosphorylates EIF4G1, which modulates EIF4G1 binding to MKNK1 and may be involved in the regulation of EIF4E phosphorylation. Phosphorylates KIT, leading to inhibition of KIT activity. Phosphorylates ATF2 which promotes cooperation between ATF2 and JUN, activating transcription. Phosphorylates SOCS2 at 'Ser- 52' facilitating its ubiquitination and proteasomal degradation (PubMed:[31578312](#)). Phosphorylates KLHL3 in response to angiotensin II signaling, decreasing the interaction between KLHL3 and WNK4 (By similarity). Phosphorylates and activates LRRK1, which phosphorylates RAB proteins involved in intracellular trafficking (By similarity).

## Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Mitochondrion membrane {ECO:0000250|UniProtKB:P17252}; Peripheral membrane protein {ECO:0000250|UniProtKB:P17252}. Nucleus Note=Translocated to the cell periphery upon tetradecanoyl phorbol acetate (TPA) treatment


## Background

This is a calcium-activated, phospholipid-dependent, serine-and threonine-specific enzyme. May play a role in cell motility by phosphorylating CSPG4 (By similarity). PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters.


## References

Yang, L., et al. Biochem. Biophys. Res. Commun. 400(1):16-20(2010)  
Jin, K., et al. J. Neurosci. 30(36):11902-11916(2010)  
Peng, H., et al. Pflugers Arch. 460(4):791-802(2010)  
Wu, M., et al. Dev. Cell 19(1):114-125(2010)  
Debata, P.R., et al. Biochem. Biophys. Res. Commun. 397(3):401-406(2010)

## Images





Mouse Prkca Antibody (N-term) (Cat. #AP14277a) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the Prkca antibody detected the Prkca protein (arrow).



Mouse Prkca Antibody (N-term) (Cat. #AP14277a) western blot analysis in mouse Neuro-2a cell line lysates (35ug/lane). This demonstrates the Prkca antibody detected the Prkca protein (arrow).

Mouse Prkca Antibody (N-term) (Cat. #AP14277a) flow cytometric analysis of Neuro-2a cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.



Mouse Prkca Antibody (N-term)  
(AP14277a) immunohistochemistry analysis in formalin fixed and paraffin embedded mouse brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Mouse Prkca Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.