

# OTU7B Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14281b

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">Q6GQQ9</a>
<b>Other Accession</b>	<a href="#">B2RUR8</a> , <a href="#">NP_064590.2</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB34874
<b>Calculated MW</b>	92526
<b>Antigen Region</b>	757-784

## Additional Information

---

<b>Gene ID</b>	56957
<b>Other Names</b>	OTU domain-containing protein 7B, Cellular zinc finger anti-NF-kappa-B protein, Zinc finger A20 domain-containing protein 1, Zinc finger protein Cezanne, OTUD7B, ZA20D1
<b>Target/Specificity</b>	This OTU7B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 757-784 amino acids from the C-terminal region of human OTU7B.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	OTU7B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	OTUD7B
<b>Synonyms</b>	ZA20D1

<b>Function</b>	Negative regulator of the non-canonical NF-kappa-B pathway that acts by mediating deubiquitination of TRAF3, an inhibitor of the NF-kappa-B pathway, thereby acting as a negative regulator of B-cell responses (PubMed: <a href="#">18178551</a> ). In response to non-canonical NF-kappa-B stimuli, deubiquitinates 'Lys-48'-linked polyubiquitin chains of TRAF3, preventing TRAF3 proteolysis and over-activation of non-canonical NF- kappa-B (By similarity). Negatively regulates mucosal immunity against infections (By similarity). Deubiquitinates ZAP70, and thereby regulates T cell receptor (TCR) signaling that leads to the activation of NF-kappa-B (PubMed: <a href="#">26903241</a> ). Plays a role in T cell homeostasis and is required for normal T cell responses, including production of IFNG and IL2 (By similarity). Mediates deubiquitination of EGFR (PubMed: <a href="#">22179831</a> ). Has deubiquitinating activity toward 'Lys-11', 'Lys- 48' and 'Lys-63'-linked polyubiquitin chains (PubMed: <a href="#">11463333</a> , PubMed: <a href="#">20622874</a> , PubMed: <a href="#">23827681</a> , PubMed: <a href="#">27732584</a> ). Has a much higher catalytic rate with 'Lys-11'-linked polyubiquitin chains (in vitro); however the physiological significance of these data are unsure (PubMed: <a href="#">27732584</a> ). Hydrolyzes both linear and branched forms of polyubiquitin (PubMed: <a href="#">12682062</a> ). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked deubiquitination of MLST8, thereby promoting assembly of the mTORC2 complex, while inhibiting formation of the mTORC1 complex (PubMed: <a href="#">28489822</a> ).
<b>Cellular Location</b>	Cytoplasm. Nucleus Note=Shuttles be cytoplasm and the nucleus in a XPO1/CRM1-dependent manner.
<b>Tissue Location</b>	Widely expressed. Abundant in kidney, heart and fetal liver. Expressed differentially among B-cells at distinct developmental stages. Higher expression seen in primary immature B- cells as compared to the mature cells.

## Background

---

OTU7B has deubiquitinating activity that is directed towards 'Lys-48' or 'Lys-63'-linked polyubiquitin chains. Hydrolyzes both linear and branched forms of polyubiquitin. Negative regulator of nuclear factor NF-kappa-B.

## References



---

Bremm, A., et al. Nat. Struct. Mol. Biol. 17(8):939-947(2010)  
Jin, Z., et al. Cell 137(4):721-735(2009)  
Enesa, K., et al. J. Biol. Chem. 283(27):18582-18590(2008)  
Bohgaki, M., et al. Biochim. Biophys. Acta 1783(5):826-837(2008)  
Enesa, K., et al. J. Biol. Chem. 283(11):7036-7045(2008)

## Images

---

OTU7B Antibody (C-term) (Cat. #AP14281b) western blot analysis in MDA-MB453 cell line lysates (35ug/lane).This demonstrates the OTU7B antibody detected the OTU7B protein (arrow).



OTU7B Antibody (C-term)  
(AP14281b) immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of OTU7B Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.