

LLGL1/2 Antibody (Center S650/654)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP14311c

Product Information

Application WB, E	
Primary Accession <u>Q6P1M3</u>	
Other Accession Q3TJ91, Q8K4K5, Q80Y17, Q15334, Q8MKF0, NP_0045	<u>4515.2</u> , <u>NP_001015002.1</u> ,
<u>NP_001026973.1</u>	
Reactivity Human	
Predicted Bovine, Mouse, Rat	
Host Rabbit	
Clonality Polyclonal	
Isotype Rabbit IgG	
Clone Names RB07627	
Calculated MW 113448	
Antigen Region 624-653	

Additional Information

Gene ID	3993
Other Names	Lethal(2) giant larvae protein homolog 2, HGL, LLGL2
Target/Specificity	This LLGL1/2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 624-653 amino acids from the Central region of human LLGL1/2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	LLGL1/2 Antibody (Center S650/654) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	LLGL2
Function	Part of a complex with GPSM2/LGN, PRKCI/aPKC and PARD6B/Par- 6, which may ensure the correct organization and orientation of bipolar spindles for

normal cell division. This complex plays roles in the initial phase of the
establishment of epithelial cell polarity.Cellular LocationCytoplasm. Note=Localized in the perinuclear structure and faintly at the cell-
cell contacts sites in the interphase. Localized at the cell periphery during
metaphase. Cortical localization in mitotic cells. Found in the lateral region of
polarized epithelial cells

Background

The lethal (2) giant larvae protein of Drosophila plays a role in asymmetric cell division, epithelial cell polarity, and cell migration. This human gene encodes a protein similar to lethal (2) giant larvae of Drosophila. In fly, the protein's ability to localize cell fate determinants is regulated by the atypical protein kinase C (aPKC). In human, this protein interacts with aPKC-containing complexes and is cortically localized in mitotic cells. Alternative splicing results in multiple transcript variants encoding different isoforms.

References

Lisovsky, M., et al. Hum. Pathol. 41(6):902-909(2010) Yoshida, T., et al. Int. J. Mol. Med. 25(4):649-656(2010) Oguri, M., et al. Am. J. Hypertens. 23(1):70-77(2010) Yamada, Y., et al. Atherosclerosis 207(1):144-149(2009) Lisovsky, M., et al. Mod. Pathol. 22(7):977-984(2009)

Images

HL-60 250 130 - ◀	LLGL1/2 Antibody (pS650/654) (Cat. #AP14311c) western blot analysis in HL-60 cell line lysates (35ug/lane).This demonstrates the LLGL1/2 antibody detected the LLGL1/2 protein (arrow).
95 72 -	
55	

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.