10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PERP Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14508b

Product Information

Application	WB, IHC-P, E
Primary Accession	Q96FX8
Other Accession	NP_071404.2
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB34260
Calculated MW	21386
Antigen Region	162-191

Additional Information

Gene ID	64065
Other Names	p53 apoptosis effector related to PMP-22, Keratinocyte-associated protein 1, KCP-1, P53-induced protein PIGPC1, Transmembrane protein THW, PERP, KCP1, KRTCAP1, PIGPC1, THW
Target/Specificity	This PERP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 162-191 amino acids from the C-terminal region of human PERP.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PERP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PERP (HGNC:17637)
Function	Component of intercellular desmosome junctions (By similarity). Plays a

	role in stratified epithelial integrity and cell- cell adhesion by promoting desmosome assembly (By similarity). Thereby plays a role in barrier function of the skin against infection (By similarity). Plays a role in mammary epithelial tissue homeostasis and remodeling during and after pregnancy, potentially via its involvement in desmosome cell-cell junctions (By similarity). Required for tooth enamel development via facilitating desmosome-mediated ameloblast adhesion to the stratum intermedium during the transitional stage of amelogenesis (By similarity). May also play a role in downstream transcriptional regulation of other genes involved in amelogenesis such as AMBN, ENAM, MMP20 and KLK4 (By similarity). Plays a role as an effector in the TP53-dependent apoptotic pathway (By similarity). Positively regulates apoptosis in T-helper 17 (Th17) cell populations via caspase-dependent signaling (By similarity). Promotes neutrophil transepithelial migration in response to chemoattractants such as hepoxilin A3 (HXA3), N-Formylmethionyl-leucyl-phenylalanine (fMLP) and CXCL8/IL-8 (PubMed:25486861). Required for neutrophil transepithelial migration in response to S.typhimurium infection (PubMed:25486861). May act as a positive regulator of endothelial cell apoptosis in response to blood flow-derived shear stress (By similarity).
Cellular Location	Cell junction, desmosome {ECO:0000250|UniProtKB:Q9JK95}. Cell membrane; Multi-pass membrane protein. Cytoplasm. Note=Associated with desmosomes (By similarity). Colocalizes with KRT14 in the cell membrane (PubMed:31898316). Clusters in a punctate pattern throughout the epithelial cytoplasm, in response to S.typhimurium infection (PubMed:25486861). {ECO:0000250|UniProtKB:Q9JK95, ECO:0000269|PubMed:25486861, ECO:0000269|PubMed:31898316}
Tissue Location	Expressed in skin, heart, placental, liver, pancreas, keratinocytes and dermal fibroblasts. May translocate to the intestinal apical epithelial cell surface via sipA and sctB1/sipC- promoted exocytic translocation following infection by S. Typhimurium (PubMed:25486861, PubMed:27078059).

Background


Component of intercellular desmosome junctions. Plays a role in stratified epithelial integrity and cell-cell adhesion by promoting desmosome assembly. Plays a role as an effector in the TP53-dependent apoptotic pathway (By similarity).


References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)
Yarden, R.I., et al. Mol. Carcinog. 49(6):545-555(2010)
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Beaudry, V.G., et al. Am. J. Med. Genet. A 149A (9), 1952-1957 (2009) :

Images

PERP Antibody (C-term) (Cat. #AP14508b) western blot analysis in NCI-H292 cell line lysates (35ug/lane).This demonstrates the PERP antibody detected the PERP protein (arrow).





PERP Antibody (C-term) (AP14508b)immunohistochemistry analysis in formalin fixed and paraffin embedded human heart tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of PERP Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.