

CNOT4 Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP1462d

Product Information

Application	WB, FC, IHC-P, E
Primary Accession	O95628
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB22947
Calculated MW	63510

Additional Information

Gene ID	4850
Other Names	CCR4-NOT transcription complex subunit 4, 632-, CCR4-associated factor 4, E3 ubiquitin-protein ligase CNOT4, Potential transcriptional repressor NOT4Hp, CNOT4, NOT4
Target/Specificity	This CNOT4 antibody is generated from rabbits immunized with human CNOT4 recombinant protein.
Dilution	WB~~1:1000 FC~~1:10~50 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CNOT4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CNOT4
Synonyms	NOT4
Function	Has E3 ubiquitin ligase activity, promoting ubiquitination and degradation of target proteins (PubMed: 11823428 , PubMed: 22159038 ,

PubMed:[26575292](#)). Involved in activation of the JAK/STAT pathway (PubMed:[11823428](#), PubMed:[22159038](#)). Catalyzes ubiquitination of methylated RBM15 (PubMed:[26575292](#)). Plays a role in quality control of translation of mitochondrial outer membrane-localized mRNA (PubMed:[29861391](#)). As part of the PINK1-regulated signaling, upon mitochondria damage, ubiquitinates ABCE1 and thereby recruits autophagy receptors to the mitochondrial outer membrane to initiate mitophagy (PubMed:[29861391](#)).

Cellular Location

Cytoplasm. Nucleus.


Background

CNOT4 has intrinsic E3 ubiquitin ligase activity. Binding of the CNOT4 RING finger to the ubiquitin-conjugating enzyme (E2) UbcH5B is highly selective. The CCR4-NOT complex functions as a global regulator of RNA polymerase II transcription.


References

Albert T.K., Nucleic Acids Res. 28:809-817(2000).
Albert T.K., EMBO J. 21:355-364(2002).

Images




AP1462D staining CNOT4 in human testis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Samples were incubated with primary antibody (1/250) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing U-2 OS cells stained with AP1462d(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP1462d, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OE188374) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

Anti-CNOT4 Antibody (N-Term) at 1:2000 dilution + Mouse cerebellum lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.