

NQO2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14656a

Product Information

| | |
|-------------------|-----------------------------|
| Application | WB, E |
| Primary Accession | P16083 |
| Other Accession | NP_000895.2 |
| Reactivity | Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB33414 |
| Calculated MW | 25919 |
| Antigen Region | 51-79 |

Additional Information

| | |
|--------------------|--|
| Gene ID | 4835 |
| Other Names | Ribosyldihyronicotinamide dehydrogenase [quinone], NRH dehydrogenase [quinone] 2, NRH:quinone oxidoreductase 2, Quinone reductase 2, QR2, NQO2, NMOR2 |
| Target/Specificity | This NQO2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 51-79 amino acids from the N-terminal region of human NQO2. |
| Dilution | WB~~1:1000 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | NQO2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|----------|-------|
| Name | NQO2 |
| Synonyms | NMOR2 |

| | |
|--------------------------|--|
| Function | The enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis. |
| Cellular Location | Cytoplasm. |


Background

NQO2 (EC 1.10.99.2) is a flavoprotein that catalyzes the 2-electron reduction of various quinones, redox dyes, and the vitamin K menadione. NQO2 predominantly uses dihydronicotinamide riboside (NRH) as the electron donor (summary by Wu et al., 1997 [PubMed 9367528]).

References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
 Ji, L.D., et al. J. Pineal Res. 48(2):133-141(2010)
 Yu, K.D., et al. Breast Cancer Res. Treat. 118(3):647-649(2009)
 Choi, J.Y., et al. Clin. Cancer Res. 15(16):5258-5266(2009)
 Yu, K.D., et al. Hum. Mol. Genet. 18(13):2502-2517(2009)

Images



NQO2 Antibody (N-term) (Cat. #AP14656a) western blot analysis in mouse spleen tissue lysates (35ug/lane). This demonstrates the NQO2 antibody detected the NQO2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.