

ALDH1A1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP1465C

Product Information

Application	WB, IHC-P, IF, FC, E
Primary Accession	P00352
Other Accession	Q35945 , Q8HYE4
Reactivity	Human, Rat, Mouse
Predicted	Monkey, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	54862
Antigen Region	302-331

Additional Information

Gene ID	216
Other Names	Retinal dehydrogenase 1, RALDH 1, RaIDH1, ALDH-E1, ALHDII, Aldehyde dehydrogenase family 1 member A1, Aldehyde dehydrogenase, cytosolic, ALDH1A1, ALDC, ALDH1, PUMB1
Target/Specificity	This ALDH1A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 302-331 amino acids from the Central region of human ALDH1A1.
Dilution	WB~~1:1000 IHC-P~~1:100~500 IF~~1:10~50 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ALDH1A1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ALDH1A1 (HGNC:402)
Function	Cytosolic dehydrogenase that catalyzes the irreversible oxidation of a wide

range of aldehydes to their corresponding carboxylic acid (PubMed:[12941160](#), PubMed:[15623782](#), PubMed:[17175089](#), PubMed:[19296407](#), PubMed:[25450233](#), PubMed:[26373694](#)). Functions downstream of retinol dehydrogenases and catalyzes the oxidation of retinaldehyde into retinoic acid, the second step in the oxidation of retinol/vitamin A into retinoic acid (By similarity). This pathway is crucial to control the levels of retinol and retinoic acid, two important molecules which excess can be teratogenic and cytotoxic (By similarity). Also oxidizes aldehydes resulting from lipid peroxidation like (E)-4-hydroxynon-2-enal/HNE, malonaldehyde and hexanal that form protein adducts and are highly cytotoxic. By participating for instance to the clearance of (E)-4-hydroxynon-2-enal/HNE in the lens epithelium prevents the formation of HNE-protein adducts and lens opacification (PubMed:[12941160](#), PubMed:[15623782](#), PubMed:[19296407](#)). Also functions downstream of fructosamine-3-kinase in the fructosamine degradation pathway by catalyzing the oxidation of 3-deoxyglucosone, the carbohydrate product of fructosamine 3-phosphate decomposition, which is itself a potent glycating agent that may react with lysine and arginine side-chains of proteins (PubMed:[17175089](#)). Also has an aminobutyraldehyde dehydrogenase activity and is probably part of an alternative pathway for the biosynthesis of GABA/4-aminobutanoate in midbrain, thereby playing a role in GABAergic synaptic transmission (By similarity).

Cellular Location	Cytoplasm, cytosol. Cell projection, axon {ECO:0000250 UniProtKB:P24549}
Tissue Location	Expressed by erythrocytes (at protein level).


Background

ALDH1A1 belongs to the aldehyde dehydrogenases family of proteins. Aldehyde dehydrogenase is the second enzyme of the major oxidative pathway of alcohol metabolism. Two major liver isoforms of this enzyme, cytosolic and mitochondrial, can be distinguished by their electrophoretic mobilities, kinetic properties, and subcellular localizations. Most Caucasians have two major isozymes, while approximately 50% of Orientals have only the cytosolic isozyme, missing the mitochondrial isozyme. A remarkably higher frequency of acute alcohol intoxication among Orientals than among Caucasians could be related to the absence of the mitochondrial isozyme.

References

Moore,S., J Stud Alcohol Drugs 68 (2), 192-196 (2007)
 Collard,F., Biochimie 89 (3), 369-373 (2007)

Images



All lanes : Anti-ANGPT2 Antibody (C-term) at 1:500 dilution
 Lane 1: U-251 MG whole cell lysate
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 55kDa
 Blocking/Dilution buffer: 5% NFDm/TBST.

Citations

- [N6-methyladenosine modification of B7-H3 mRNA promotes the development and progression of colorectal cancer](#)
- [A novel reporter construct for screening small molecule inhibitors that specifically target self-renewing cancer cells](#)
- [Cargo-free nano-medicine with pH-sensitivity for co-delivery of DOX conjugated prodrug with SN38 to synergistically eradicate breast cancer stem cells](#)
- [Ataxin-1 regulates the cerebellar bioenergetics proteome through the GSK3 \$\beta\$ -mTOR pathway which is altered in Spinocerebellar ataxia type 1 \(SCA1\)](#)
- [Wnt ligands from the embryonic surface ectoderm regulate 'bimetallic strip' optic cup morphogenesis in mouse](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.