

# TCEA1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14761c

## Product Information

---

Application	WB, IHC-P, E
Primary Accession	<a href="#">P23193</a>
Other Accession	<a href="#">NP_006747.1</a> , <a href="#">NP_958845.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35010
Calculated MW	33970
Antigen Region	98-126

## Additional Information

---

Gene ID	6917
Other Names	Transcription elongation factor A protein 1, Transcription elongation factor S-II protein 1, Transcription elongation factor TFIISo, TCEA1, GTF2S, TFIIS
Target/Specificity	This TCEA1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 98-126 amino acids from the Central region of human TCEA1.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TCEA1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	TCEA1
Synonyms	GTF2S, TFIIS
Function	Necessary for efficient RNA polymerase II transcription elongation past

template-encoded arresting sites. The arresting sites in DNA have the property of trapping a certain fraction of elongating RNA polymerases that pass through, resulting in locked ternary complexes. Cleavage of the nascent transcript by S-II allows the resumption of elongation from the new 3'-terminus.

**Cellular Location**

Nucleus.

## Background

---

Necessary for efficient RNA polymerase II transcription elongation past template-encoded arresting sites. The arresting sites in DNA have the property of trapping a certain fraction of elongating RNA polymerases that pass through, resulting in locked ternary complexes. Cleavage of the nascent transcript by S-II allows the resumption of elongation from the new 3'-terminus.


## References

---


Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)  
Matsuoka, S., et al. Science 316(5828):1160-1166(2007)  
Olsen, J.V., et al. Cell 127(3):635-648(2006)  
Olsen, J.V., et al. Cell 127(3):635-648(2006)  
Asp, J., et al. Genes Chromosomes Cancer 45(9):820-828(2006)

## Images

---



TCEA1 Antibody (Center) (Cat. #AP14761c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the TCEA1 antibody detected the TCEA1 protein (arrow).



TCEA1 Antibody (Center) (AP14761c) immunohistochemistry analysis in formalin fixed and paraffin embedded human tonsil tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TCEA1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.