10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ACTA1/Alpha-actin Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14779b

Product Information

Application	IHC-P, IF, FC, WB, E
Primary Accession	P68133
Other Accession	P68136, P68135, P68137, P68134, P68139, P68138, P04751, P68035, P68033, P68032, P68034, Q3ZC07, P62738, P62740, P62737, P62736, P08023, P62739, P04752, P10995, NP_001091.1
Reactivity	Human, Rat, Mouse
Predicted	Mouse, Rat, Rabbit, Pig, Chicken, Bovine, Xenopus, Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	42051
Antigen Region	346-375

Additional Information

Gene ID	58
Other Names	Actin, alpha skeletal muscle, Alpha-actin-1, ACTA1, ACTA
Target/Specificity	This ACTA1/Alpha-actin antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 346-375 amino acids from the C-terminal region of human ACTA1/Alpha-actin.
Dilution	IHC-P~~1:100~500 IF~~1:10~50 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ACTA1/Alpha-actin Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ACTA1
Synonyms	ACTA

Function	Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.
Cellular Location	Cytoplasm, cytoskeleton.


Background

The product encoded by this gene belongs to the actin family of proteins, which are highly conserved proteins that play a role in cell motility, structure and integrity. Alpha, beta and gamma actin isoforms have been identified, with alpha actins being a major constituent of the contractile apparatus, while beta and gamma actins are involved in the regulation of cell motility. This actin is an alpha actin that is found in skeletal muscle. Mutations in this gene cause nemaline myopathy type 3, congenital myopathy with excess of thin myofilaments, congenital myopathy with cores, and congenital myopathy with fiber-type disproportion, diseases that lead to muscle fiber defects.


References

Kim, E.Y., et al. Am. J. Physiol. Renal Physiol. 299 (3), F594-F604 (2010) :
Haigh, S.E., et al. Neuromuscul. Disord. 20(6):363-374(2010)
Yu, G., et al. J Clin Neurosci 17(6):766-769(2010)
Yu, C.H., et al. PLoS ONE 5 (7), E11878 (2010) :
Licastro, F., et al. Curr. Pharm. Des. 16(7):783-788(2010)

Images





ACTA1/α-actin Antibody (C-term) (Cat. #AP14779b) western blot analysis in A549,RD cell line and human placenta lysates (35ug/lane).This demonstrates the ACTA1/α-actin antibody detected the ACTA1/α-actin protein (arrow).



ACTA1/?-actin Antibody (C-term) (AP14779b)immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of ACTA1/?-actin Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Confocal immunofluorescent analysis of ACTA1/?-actin Antibody (C-term)(Cat#AP14779b) with A549 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green).DAPI was used to stain the cell nuclear (blue).





ACTA1/?-actin Antibody (C-term) (Cat. #AP14779b) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Citations

- Exposure to concentrated ambient particulate matter induces reversible increase of heart weight in spontaneously hypertensive rats.
- Directed Differentiation of Zebrafish Pluripotent Embryonic Cells to Functional Cardiomyocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.