

ZNF398 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP14859c

Product Information

Application	WB, E
Primary Accession	Q8TD17
Other Accession	NP_065832.1 , NP_733787.1
Reactivity	Human
Predicted	Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB34998
Calculated MW	71311
Antigen Region	186-215

Additional Information

Gene ID	57541
Other Names	Zinc finger protein 398, Zinc finger DNA-binding protein p52/p71, ZNF398, KIAA1339, ZER6
Target/Specificity	This ZNF398 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 186-215 amino acids from the Central region of human ZNF398.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ZNF398 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ZNF398
Synonyms	KIAA1339, ZER6

Function	Functions as a transcriptional activator.
Cellular Location	Nucleus.


Background

This gene encodes a member of the Kruppel family of C2H2-type zinc-finger transcription factor proteins. The encoded protein acts as a transcriptional activator. Two transcript variants encoding distinct isoforms have been identified for this gene. Other transcript variants have been described, but their full length sequence has not been determined.

References

Conroy, A.T., et al. J. Biol. Chem. 277(11):9326-9334(2002)

Images



ZNF398 Antibody (Center) (Cat. #AP14859c) western blot analysis in SK-BR-3 cell line lysates (35ug/lane). This demonstrates the ZNF398 antibody detected the ZNF398 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.