

# NANOG Antibody (S285)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP1486E

## Product Information

---

Application	WB, IF, E
Primary Accession	<a href="#">Q9H9S0</a>
Other Accession	<a href="#">Q6NSW7</a> , <a href="#">NP_079141</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB18922
Calculated MW	34620
Antigen Region	267-292

## Additional Information

---

Gene ID	79923
Other Names	Homeobox protein NANOG, Homeobox transcription factor Nanog, hNanog, NANOG
Target/Specificity	This NANOG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 267-292 amino acids from human NANOG.
Dilution	WB~~1:1000 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NANOG Antibody (S285) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	NANOG
Function	Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and

trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. Binds to the POU5F1/OCT4 promoter (PubMed:[25825768](#)). Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

#### Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00108, ECO:0000269 | PubMed:15983365}

#### Tissue Location

Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.


## Background

NANOG is a Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. It imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. This protein blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. NANOG acts as a transcriptional activator or repressor. It binds optimally to the DNA consensus sequence 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. When overexpressed, this protein promotes cells to enter into S phase and proliferation.

## References



Kochupurakkal, B.S., Biochem. Biophys. Res. Commun. 365 (4), 846-850 (2008)  
Freberg, C.T., Mol. Biol. Cell 18 (5), 1543-1553 (2007)

## Images



All lanes : Anti-NANOG Antibody (S285) at 1:1000 dilution  
Lane 1: MCF-7 whole cell lysate Lane 2: human testis lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Western blot analysis of NANOG Antibody (S285) (Cat.# AP1486e) in HepG2 cell line lysates (35ug/lane). NANOG (arrow) was detected using the purified Pab.



Fluorescent confocal image of SY5Y cells stained with AP1486e NANOG (S285) antibody. SY5Y cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min), then incubated with AP1486e NANOG (S285) primary antibody (1:200, 2 h at room temperature). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (5.25  $\mu$ M, 25 min). Nuclei were counterstained with Hoechst 33342 (blue) (10  $\mu$ g/ml, 3 min). Nanog immunoreactivity is localized mainly to the nuclei and also to the cytoplasm.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.