10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SMAD9 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14903C

Product Information

Application	WB, IHC-P-Leica, E
Primary Accession	O15198
Other Accession	NP_005896.1, NP_001120689.1
Reactivity	Human
Predicted	Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35231
Calculated MW	52493
Antigen Region	200-228

Additional Information

Gene ID	4093
Other Names	Mothers against decapentaplegic homolog 9, MAD homolog 9, Mothers against DPP homolog 9, Madh6, SMAD family member 9, SMAD 9, Smad9, SMAD9, MADH6, MADH9
Target/Specificity	This SMAD9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 200-228 amino acids from the Central region of human SMAD9.
Dilution	WB~~1:1000 IHC-P-Leica~~1:500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SMAD9 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SMAD9 (HGNC:6774)

Function	Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD9 is a receptor- regulated SMAD (R-SMAD).
Cellular Location	Cytoplasm. Nucleus. Note=In the cytoplasm in the absence of ligand. Migration to the nucleus when complexed with SMAD4 (By similarity).
Tissue Location	Expressed in heart, brain, placenta, lung, skeletal muscle, prostate, testis, ovary and small intestine. Also expressed in fetal brain, lung and kidney

Background

The protein encoded by this gene is a member of the SMAD family, which transduces signals from TGF-beta family members. The encoded protein is activated by bone morphogenetic proteins and interacts with SMAD4. Two transcript variants encoding different isoforms have been found for this gene.

References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Liu, Y., et al. FASEB J. 23(7):2299-2306(2009)
Su, D., et al. J. Biol. Chem. 284(18):12153-12164(2009)
Shintani, M., et al. J. Med. Genet. 46(5):331-337(2009)
Hoover, L.L., et al. Sci Signal 1 (46), PE48 (2008) :

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.