

# VILIP3 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP1563a

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, IHC-P, E           |
| <b>Primary Accession</b> | <a href="#">P37235</a> |
| <b>Reactivity</b>        | Human, Mouse           |
| <b>Host</b>              | Rabbit                 |
| <b>Clonality</b>         | Polyclonal             |
| <b>Isotype</b>           | Rabbit IgG             |
| <b>Clone Names</b>       | RB0501                 |
| <b>Calculated MW</b>     | 22313                  |
| <b>Antigen Region</b>    | 2-32                   |

## Additional Information

---

|                           |   |
|---------------------------|---|
| <b>Gene ID</b>            | 3241  |
| <b>Other Names</b>        | Hippocalcin-like protein 1, Calcium-binding protein BDR-1, HLP2, Visinin-like protein 3, VILIP-3, HPCAL1, BDR1  |
| <b>Target/Specificity</b> | This VILIP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 2-32 amino acids from the N-terminal region of human VILIP3.                     |
| <b>Dilution</b>           | WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.   |
| <b>Format</b>             | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS. |
| <b>Storage</b>            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.   |
| <b>Precautions</b>        | VILIP3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.  |

## Protein Information

---

|                 |   |
|-----------------|---|
| <b>Name</b>     | HPCAL1  |
| <b>Synonyms</b> | BDR1  |
| <b>Function</b> | May be involved in the calcium-dependent regulation of rhodopsin phosphorylation. |


## Background

VILIP3 is a member of neuron-specific calcium-binding proteins family found in the retina and brain. It is highly similar to human hippocalcin protein and nearly identical to the rat and mouse hippocalcin like-1 proteins. It may be involved in the calcium-dependent regulation of rhodopsin phosphorylation and may be of relevance for neuronal signalling in the central nervous system. There are two alternatively spliced transcript variants of this gene, with multiple polyadenylation sites. Transcript variant 1 utilizes a different exon and also lacks one exon in the 5' UTR, as compared to variant 2; thus, the encoded protein is the same.


## References

Braunewell, K., et al., Dement Geriatr Cogn Disord 12(2):110-116 (2001). Bernstein, H.G., et al., J Neurocytol 28(8):655-662 (1999). Kobayashi, M., et al., Biochim. Biophys. Acta 1222(3):515-518 (1994). Hidaka, H., et al., Neurosci. Res. 16(2):73-77 (1993). Ivings, L., et al., Biochem. J. 363 (Pt 3), 599-608 (2002).


## Images



Western blot analysis of VILIP3 (arrow) using rabbit polyclonal VILIP3 Antibody (N-term) (Cat.#AP1563a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the VILIP3 gene (Lane 2) (Origene Technologies).



The anti-VILIP3 Pab (Cat. #AP1563a) is used in Western blot to detect VILIP3 in mouse cerebellum tissue lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.