

SOX11 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP16147a

Product Information

Application	WB, E
Primary Accession	P35716
Other Accession	P48435 , NP_003099.1
Reactivity	Human, Rat, Mouse
Predicted	Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35379
Calculated MW	46679
Antigen Region	103-132

Additional Information

Gene ID	6664
Other Names	Transcription factor SOX-11, SOX11
Target/Specificity	This SOX11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 103-132 amino acids from the N-terminal region of human SOX11.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SOX11 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SOX11
Function	Transcription factor that acts as a transcriptional activator (PubMed: 24886874 , PubMed: 26543203). Binds cooperatively with POU3F2/BRN2 or POU3F1/OCT6 to gene promoters, which enhances

transcriptional activation (By similarity). Acts as a transcriptional activator of TEAD2 by binding to its gene promoter and first intron (By similarity). Plays a redundant role with SOX4 and SOX12 in cell survival of developing tissues such as the neural tube, branchial arches and somites, thereby contributing to organogenesis (By similarity).

Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00267, ECO:0000269 | PubMed:24886874, ECO:0000269 | PubMed:35938035}

Tissue Location

Expressed primarily in the brain and heart, with low expression in the kidney, pancreas and muscle


Background

This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. The encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins. The protein may function in the developing nervous system and play a role in tumorigenesis.


References

Kottgen, A., et al. Nat. Genet. 42(5):376-384(2010)
Fernandez, V., et al. Cancer Res. 70(4):1408-1418(2010)
Dictor, M., et al. Haematologica 94(11):1563-1568(2009)
Mozos, A., et al. Haematologica 94(11):1555-1562(2009)
Hide, T., et al. Cancer Res. 69(20):7953-7959(2009)

Images



All lanes : Anti-SOX11 Antibody (N-term) at 1:2000 dilution
Lane 1: rat brain lysates
Lane 2: human brain lysates
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution
Predicted band size : 47 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



SOX11 Antibody (N-term) (Cat. #AP16147a) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the SOX11 antibody detected the SOX11 protein (arrow).