

RPL12 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP16275c

Product Information

Application	WB, E
Primary Accession	P30050
Other Accession	P23358 , P35979 , P61284 , NP_000967.1
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35573
Calculated MW	17819
Antigen Region	71-100

Additional Information

Gene ID	6136
Other Names	60S ribosomal protein L12, RPL12
Target/Specificity	This RPL12 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 71-100 amino acids from the Central region of human RPL12.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RPL12 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RPL12
Function	Component of the large ribosomal subunit (PubMed: 25901680). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed: 25901680). Binds directly to 26S ribosomal

RNA (PubMed:[25901680](#)).

Cellular Location

Cytoplasm


Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L11P family of ribosomal proteins. It is located in the cytoplasm. The protein binds directly to the 26S rRNA. This gene is co-transcribed with the U65 snoRNA, which is located in its fourth intron. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

References

Letra, A., et al. Am. J. Med. Genet. A 152A (7), 1701-1710 (2010) :
Wheeler, H.E., et al. PLoS Genet. 5 (10), E1000685 (2009) :
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)

Images



All lanes : Anti-RPL12 Antibody (Center) at 1:1000 dilution
Lane 1: Raji whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: mouse pancreas lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size : 19 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

- [Ribosomal Stalk Protein Silencing Partially Corrects the AF508-CFTR Functional Expression Defect.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.