

# RNF4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP16278b

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P78317</a>
<b>Other Accession</b>	<a href="#">NP_001171939.1</a> , <a href="#">NP_001171938.1</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	21319
<b>Antigen Region</b>	95-123

## Additional Information

---

<b>Gene ID</b>	6047
<b>Other Names</b>	E3 ubiquitin-protein ligase RNF4, 632-, RING finger protein 4, Small nuclear ring finger protein, Protein SNURF, RNF4, SNURF
<b>Target/Specificity</b>	This RNF4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 95-123 amino acids from the C-terminal region of human RNF4.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:250 FC~~1:25 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	RNF4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	RNF4 {ECO:0000303   PubMed:15815621, ECO:0000312   HGNC:HGNC:10067}
<b>Function</b>	E3 ubiquitin-protein ligase which binds polysumoylated chains covalently attached to proteins and mediates 'Lys-6'-, 'Lys-11'-, 'Lys- 48'- and 'Lys-63'-linked polyubiquitination of those substrates and their subsequent

targeting to the proteasome for degradation (PubMed:[18408734](#), PubMed:[19307308](#), PubMed:[35013556](#)). Regulates the degradation of several proteins including PML and the transcriptional activator PEA3 (PubMed:[18408734](#), PubMed:[19307308](#), PubMed:[20943951](#)). Involved in chromosome alignment and spindle assembly, it regulates the kinetochore CENPH-CENPI-CENPK complex by targeting polysumoylated CENPI to proteasomal degradation (PubMed:[20212317](#)). Regulates the cellular responses to hypoxia and heat shock through degradation of respectively EPAS1 and PARP1 (PubMed:[19779455](#), PubMed:[20026589](#)). Alternatively, it may also bind DNA/nucleosomes and have a more direct role in the regulation of transcription for instance enhancing basal transcription and steroid receptor-mediated transcriptional activation (PubMed:[12885770](#)). Catalyzes ubiquitination of sumoylated PARP1 in response to PARP1 trapping to chromatin, leading to PARP1 removal from chromatin by VCP/p97 (PubMed:[35013556](#)).

**Cellular Location** Cytoplasm. Nucleus. Nucleus, PML body

**Tissue Location** Widely expressed at low levels in many tissues; highly expressed in testis.

## Background

---

The protein encoded by this gene contains a RING finger motif and acts as a transcription regulator. This protein has been shown to interact with, and inhibit the activity of, TRPS1, a transcription suppressor of GATA-mediated transcription. Transcription repressor ZNF278/PATZ is found to interact with this protein, and thus reduce the enhancement of androgen receptor-dependent transcription mediated by this protein. Studies of the mouse and rat counterparts suggested a role of this protein in spermatogenesis. A pseudogene of this gene is found on chromosome 1.


## References

---

- Hu, X.V., et al. Proc. Natl. Acad. Sci. U.S.A. 107(34):15087-15092(2010)  
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
Salonen, J., et al. Mol. Cell. Endocrinol. 307 (1-2), 205-210 (2009) :  
Percherancier, Y., et al. J. Biol. Chem. 284(24):16595-16608(2009)  
Tatham, M.H., et al. Nat. Cell Biol. 10(5):538-546(2008)


## Images

---




AP16278b staining RNF4 in human heart tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Samples were incubated with primary antibody (1/250) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

Overlay histogram showing HepG2 cells stained with AP16278b(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in



2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP16278b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.



Anti-RNF4 Antibody (C-term) at 1:2000 dilution + Human kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

## Citations

- [Targeting HDAC3 to overcome the resistance to ATRA or arsenic in acute promyelocytic leukemia through ubiquitination and degradation of PML-RAR \$\alpha\$](#)
- [SEN3 regulates the global protein turnover and the Sp1 level via antagonizing SUMO2/3-targeted ubiquitination and degradation.](#)
- [TRIB3 Promotes APL Progression through Stabilization of the Oncoprotein PML-RAR \$\alpha\$  and Inhibition of p53-Mediated Senescence.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.