10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BAG4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16332b

Product Information

Application	WB, E
Primary Accession	O95429
Other Accession	Q8CI61, NP_004865.1
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35817
Calculated MW	49594
Antigen Region	381-409

Additional Information

Gene ID	9530
Other Names	BAG family molecular chaperone regulator 4, BAG-4, Bcl-2-associated athanogene 4, Silencer of death domains, BAG4, SODD
Target/Specificity	This BAG4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 381-409 amino acids from the C-terminal region of human BAG4.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	BAG4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BAG4
Synonyms	SODD

Function	Inhibits the chaperone activity of HSP70/HSC70 by promoting substrate release (By similarity). Prevents constitutive TNFRSF1A signaling. Negative regulator of PRKN translocation to damaged mitochondria.
Cellular Location	Cytoplasm.
Tissue Location	Ubiquitous.


Background

BAG4 is a member of the BAG1-related protein family. BAG1 is an anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors, growth factor receptors and members of the heat shock protein 70 kDa family. This protein contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone activity of Hsc70/Hsp70. This protein was found to be associated with the death domain of tumor necrosis factor receptor type 1 (TNF-R1) and death receptor-3 (DR3), and thereby negatively regulates downstream cell death signaling. The regulatory role of this protein in cell death was demonstrated in epithelial cells which undergo apoptosis while integrin mediated matrix contacts are lost.

References

Bailey, S.D., et al. Diabetes Care (2010) In press :
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Tao, H.F., et al. Zhongguo Shi Yan Xue Ye Xue Za Zhi 15(3):501-505(2007)
Riley, B.M., et al. Am. J. Med. Genet. A 143A (8), 846-852 (2007) :
Yang, Z.Q., et al. Cancer Res. 66(24):11632-11643(2006)

Images



BAG4 Antibody (C-term) (Cat. #AP16332b) western blot analysis in Hela cell line lysates (35ug/lane).This demonstrates the BAG4 antibody detected the BAG4 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.