

EIF3C Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP16361c

Product Information

Application	WB, E
Primary Accession	Q99613
Other Accession	B5ME19 , Q4QR58 , B5DFC8 , Q8R1B4 , Q3SYW6 , NP_001032897.1 , NP_003743.1
Reactivity	Mouse
Predicted	Bovine, Rat, Xenopus, Human, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35278
Calculated MW	105344
Antigen Region	538-566

Additional Information

Gene ID	8663
Other Names	Eukaryotic translation initiation factor 3 subunit C {ECO:0000255 HAMAP-Rule:MF_03002}, eIF3c {ECO:0000255 HAMAP-Rule:MF_03002}, Eukaryotic translation initiation factor 3 subunit 8 {ECO:0000255 HAMAP-Rule:MF_03002}, eIF3 p110 {ECO:0000255 HAMAP-Rule:MF_03002}, EIF3C {ECO:0000255 HAMAP-Rule:MF_03002}
Target/Specificity	This EIF3C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 538-566 amino acids from the Central region of human EIF3C.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EIF3C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EIF3C {ECO:0000255 HAMAP-Rule:MF_03002}
Function	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed: 17581632 , PubMed: 25849773 , PubMed: 27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNA ⁱ and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed: 17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed: 25849773).
Cellular Location	Cytoplasm {ECO:0000255 HAMAP-Rule:MF_03002}.


Background

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAⁱ and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.

References

Imielinski, M., et al. Nat. Genet. 41(12):1335-1340(2009)
Zhou, M., et al. Proc. Natl. Acad. Sci. U.S.A. 105(47):18139-18144(2008)
Masutani, M., et al. EMBO J. 26(14):3373-3383(2007)
Damoc, E., et al. Mol. Cell Proteomics 6(7):1135-1146(2007)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)

Images



EIF3C Antibody (Center) (Cat. #AP16361c) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the EIF3C antibody detected the EIF3C protein (arrow).