

DNMT3L Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP16566b

Product Information

Application	WB, E
Primary Accession	Q9UJW3
Other Accession	NP_037501.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB36420
Calculated MW	43583
Antigen Region	355-381

Additional Information

Gene ID	29947
Other Names	DNA (cytosine-5)-methyltransferase 3-like, DNMT3L
Target/Specificity	This DNMT3L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 355-381 amino acids from the C-terminal region of human DNMT3L.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DNMT3L Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DNMT3L
Function	Catalytically inactive regulatory factor of DNA methyltransferases that can either promote or inhibit DNA methylation depending on the context (By similarity). Essential for the function of DNMT3A and DNMT3B: activates DNMT3A and DNMT3B by binding to their catalytic domain

(PubMed:[17687327](#)). Acts by accelerating the binding of DNA and S-adenosyl-L-methionine (AdoMet) to the methyltransferases and dissociates from the complex after DNA binding to the methyltransferases (PubMed:[17687327](#)). Recognizes unmethylated histone H3 lysine 4 (H3K4me0) and induces de novo DNA methylation by recruitment or activation of DNMT3 (PubMed:[17687327](#)). Plays a key role in embryonic stem cells and germ cells (By similarity). In germ cells, required for the methylation of imprinted loci together with DNMT3A (By similarity). In male germ cells, specifically required to methylate retrotransposons, preventing their mobilization (By similarity). Plays a key role in embryonic stem cells (ESCs) by acting both as an positive and negative regulator of DNA methylation (By similarity). While it promotes DNA methylation of housekeeping genes together with DNMT3A and DNMT3B, it also acts as an inhibitor of DNA methylation at the promoter of bivalent genes (By similarity). Interacts with the EZH2 component of the PRC2/EED-EZH2 complex, preventing interaction of DNMT3A and DNMT3B with the PRC2/EED-EZH2 complex, leading to maintain low methylation levels at the promoters of bivalent genes (By similarity). Promotes differentiation of ESCs into primordial germ cells by inhibiting DNA methylation at the promoter of RHOX5, thereby activating its expression (By similarity).

Cellular Location

Nucleus.

Tissue Location

Expressed at low levels in several tissues including testis, ovary, and thymus.

Background


CpG methylation is an epigenetic modification that is important for embryonic development, imprinting, and X-chromosome inactivation. Studies in mice have demonstrated that DNA methylation is required for mammalian development. This gene encodes a nuclear protein with similarity to DNA methyltransferases. This protein is not thought to function as a DNA methyltransferase as it does not contain the amino acid residues necessary for methyltransferase activity. However, this protein does stimulate de novo methylation by DNA cytosine methyltransferase 3 alpha and it is thought to be required for the establishment of maternal genomic imprints. This protein also mediates transcriptional repression through interaction with histone deacetylase 1. Alternative splicing results in two transcript variants. An additional splice variant has been described but its biological validity has not been determined.

References

Holz-Schietinger, C., et al. J. Biol. Chem. 285(38):29091-29100(2010)
 Manderwad, G.P., et al. Arch. Pathol. Lab. Med. 134(8):1193-1196(2010)
 Kim, H., et al. Int. J. Oncol. 36(6):1563-1572(2010)
 Minami, K., et al. Clin. Cancer Res. 16(10):2751-2759(2010)
 Haggarty, P., et al. PLoS ONE 5 (6), E11329 (2010) :

Images

Anti-DNMT3L Antibody (C-term) at 1:1000 dilution +
 Jurkat whole cell lysate Lysates/proteins at 20 µg per lane.
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase
 conjugated at 1/10000 dilution. Predicted band size : 44
 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.