

# TNNT2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16580c

## **Product Information**

Application WB, E Primary Accession P45379

Other Accession P09741, P50752, NP 001001431.1, NP 001001430.1

**Reactivity** Human, Mouse

Predicted Rabbit
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB33602
Calculated MW 35924
Antigen Region 182-211

## **Additional Information**

**Gene ID** 7139

Other Names Troponin T, cardiac muscle, TnTc, Cardiac muscle troponin T, cTnT, TNNT2

**Target/Specificity** This TNNT2 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 182-211 amino acids from the Central

region of human TNNT2.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**TNNT2 Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

### **Protein Information**

Name TNNT2

**Function** Troponin T is the tropomyosin-binding subunit of troponin, the thin

filament regulatory complex which confers calcium-sensitivity to striated

muscle actomyosin ATPase activity.

#### **Tissue Location**

Heart. The fetal heart shows a greater expression in the atrium than in the ventricle, while the adult heart shows a greater expression in the ventricle than in the atrium. Isoform 6 predominates in normal adult heart. Isoforms 1, 7 and 8 are expressed in fetal heart. Isoform 7 is also expressed in failing adult heart


# **Background**

The protein encoded by this gene is the tropomyosin-binding subunit of the troponin complex, which is located on the thin filament of striated muscles and regulates muscle contraction in response to alterations in intracellular calcium ion concentration. Mutations in this gene have been associated with familial hypertrophic cardiomyopathy as well as with dilated cardiomyopathy. Transcripts for this gene undergo alternative splicing that results in many tissue-specific isoforms, however, the full-length nature of some of these variants has not yet been determined.

## References

Millat, G., et al. Clin. Chim. Acta 411 (23-24), 1983-1991 (2010): Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Millat, G., et al. Eur J Med Genet 53(5):261-267(2010)
Watt, K.D., et al. Liver Transpl. 16(8):990-998(2010)
Koide, K., et al. Heart Vessels 25(3):217-222(2010)

## **Images**



TNNT2 Antibody (Center) (Cat. #AP16580c) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the TNNT2 antibody detected the TNNT2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.