

MED16 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP16692c

Product Information

Application	WB, E
Primary Accession	Q9Y2X0
Other Accession	NP_005472.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35935
Calculated MW	96793
Antigen Region	253-281

Additional Information

Gene ID	10025
Other Names	Mediator of RNA polymerase II transcription subunit 16, Mediator complex subunit 16, Thyroid hormone receptor-associated protein 5, Thyroid hormone receptor-associated protein complex 95 kDa component, Trap95, Vitamin D3 receptor-interacting protein complex 92 kDa component, DRIP92, MED16, DRIP92 {ECO:0000312 EMBL:AAD310871}, THRAP5
Target/Specificity	This MED16 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 253-281 amino acids from the Central region of human MED16.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MED16 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MED16
-------------	-------

Synonyms

DRIP92 {ECO:0000312 | EMBL:AAD31087.1}, TH

Function

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Cellular Location

Nucleus.

Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

References

Sato, S., et al. Mol. Cell 14(5):685-691(2004)


Tomomori-Sato, C., et al. J. Biol. Chem. 279(7):5846-5851(2004)

Wang, Q., et al. J. Biol. Chem. 277(45):42852-42858(2002)

Kang, Y.K., et al. Proc. Natl. Acad. Sci. U.S.A. 99(5):2642-2647(2002)

Suzuki, Y., et al. Genome Res. 11(5):677-684(2001)

Images



MED16 Antibody (Center) (Cat. #AP16692c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the MED16 antibody detected the MED16 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.