

PPP1CC Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP16979b

Product Information

Application	WB, E
Primary Accession	P36873
Other Accession	P63088 , P63087 , P61287 , NP_002701.1
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB36668
Calculated MW	36984
Antigen Region	281-309

Additional Information

Gene ID	5501
Other Names	Serine/threonine-protein phosphatase PP1-gamma catalytic subunit, PP-1G, Protein phosphatase 1C catalytic subunit, PPP1CC
Target/Specificity	This PPP1CC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 281-309 amino acids from the C-terminal region of human PPP1CC.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PPP1CC Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PPP1CC
Function	Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of

biological targets (PubMed:[17936702](#), PubMed:[25012651](#)). Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Dephosphorylates RPS6KB1 (PubMed:[17936702](#)). Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase (PubMed:[20516061](#)). In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation (PubMed:[21712997](#)). May dephosphorylate CSNK1D and CSNK1E (By similarity). Regulates the recruitment of the SKA complex to kinetochores (PubMed:[28982702](#)). Dephosphorylates the 'Ser-418' residue of FOXP3 in regulatory T-cells (Treg) from patients with rheumatoid arthritis, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:[23396208](#)). Together with PPP1CA (PP1- alpha subunit), dephosphorylates IFIH1/MDA5 and RIG-I leading to their activation and a functional innate immune response (PubMed:[23499489](#)). Core component of the SHOC2-MRAS-PP1c (SMP) holophosphatase complex that regulates the MAPK pathway activation (PubMed:[35768504](#), PubMed:[35831509](#)). The SMP complex specifically dephosphorylates the inhibitory phosphorylation at 'Ser-259' of RAF1 kinase, 'Ser-365' of BRAF kinase and 'Ser-214' of ARAF kinase, stimulating their kinase activities (PubMed:[35768504](#), PubMed:[35831509](#)). Dephosphorylates MKI67 at the onset of anaphase (PubMed:[25012651](#)). The SMP complex enhances the dephosphorylation activity and substrate specificity of PP1c (PubMed:[35768504](#), PubMed:[35831509](#)).

Cellular Location

Cytoplasm. Nucleus. Nucleus, nucleolus. Nucleus, nucleoplasm. Nucleus speckle. Chromosome, centromere, kinetochore. Cleavage furrow. Midbody Mitochondrion. Cytoplasm, cytoskeleton, microtubule organizing center Note=Colocalizes with SPZ1 in the nucleus (By similarity). Colocalizes with URI1 at mitochondrion (PubMed:[17936702](#)). Rapidly exchanges between the nucleolar, nucleoplasmic and cytoplasmic compartments (PubMed:[11739654](#)). Highly mobile in cells and can be relocalized through interaction with targeting subunits (PubMed:[17965019](#)). In the presence of PPP1R8 relocalizes from the nucleolus to nuclear speckles (PubMed:[11739654](#)). Shows a dynamic targeting to specific sites throughout the cell cycle (PubMed:[12529430](#)). Highly concentrated in nucleoli of interphase cells and localizes at kinetochores early in mitosis (PubMed:[12529430](#)). Relocalization to chromosome-containing regions occurs at the transition from early to late anaphase (PubMed:[12529430](#)). Also accumulates at the cleavage furrow and midbody by telophase (PubMed:[12529430](#)). Colocalizes with DYNLT4 in the microtubule organizing center (MTOC) (PubMed:[23789093](#)) {ECO:0000250|UniProtKB:P63087, ECO:0000269|PubMed:[11739654](#), ECO:0000269|PubMed:[12529430](#), ECO:0000269|PubMed:[17936702](#), ECO:0000269|PubMed:[17965019](#), ECO:0000269|PubMed:[23789093](#)}


Background

Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II.


References

Lee, J.H., et al. J. Biol. Chem. 285(32):24466-24476(2010)
Kuzmin, A., et al. Biol. Reprod. 81(2):319-326(2009)
Fujiki, R., et al. Nature 459(7245):455-459(2009)
Rogne, M., et al. Hum. Mol. Genet. 18(5):978-987(2009)
Tchivilev, I., et al. J. Biol. Chem. 283(32):22193-22205(2008)

Images



PPP1CC Antibody (C-term) (Cat. #AP16979b) western blot analysis in WiDr cell line lysates (35ug/lane). This demonstrates the PPP1CC antibody detected the PPP1CC protein (arrow).



Western blot analysis of PPP1CC (arrow) using rabbit polyclonal PPP1CC Antibody (C-term) (Cat. #AP16979b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the PPP1CC gene.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.