

MAP3K7IP3 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP17071a

Product Information

Application	WB, E
Primary Accession	Q8N5C8
Other Accession	Q571K4 , NP_690000.2
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB36935
Calculated MW	78653
Antigen Region	1-30

Additional Information

Gene ID	257397
Other Names	TGF-beta-activated kinase 1 and MAP3K7-binding protein 3, Mitogen-activated protein kinase kinase kinase 7-interacting protein 3, NF-kappa-B-activating protein 1, TAK1-binding protein 3, TAB-3, TGF-beta-activated kinase 1-binding protein 3, TAB3, MAP3K7IP3
Target/Specificity	This MAP3K7IP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human MAP3K7IP3.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MAP3K7IP3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TAB3 {ECO:0000303 PubMed:14633987, ECO:0000312 HGNC:HGNC:30681}
------	---

Function	Adapter required to activate the JNK and NF-kappa-B signaling pathways through the specific recognition of 'Lys-63'-linked polyubiquitin chains by its RanBP2-type zinc finger (NZF) (PubMed: 14633987 , PubMed: 14766965 , PubMed: 15327770 , PubMed: 22158122). Acts as an adapter linking MAP3K7/TAK1 and TRAF6 to 'Lys-63'-linked polyubiquitin chains (PubMed: 14633987 , PubMed: 14766965 , PubMed: 15327770 , PubMed: 22158122 , PubMed: 36593296). The RanBP2-type zinc finger (NZF) specifically recognizes Lys-63'-linked polyubiquitin chains unanchored or anchored to the substrate proteins such as RIPK1/RIP1 and RIPK2: this acts as a scaffold to organize a large signaling complex to promote autophosphorylation of MAP3K7/TAK1, and subsequent activation of I-kappa-B-kinase (IKK) core complex by MAP3K7/TAK1 (PubMed: 15327770 , PubMed: 18079694 , PubMed: 22158122).
Tissue Location	Widely expressed. Constitutively overexpressed in certain tumor tissues. [Isoform 2]: Minor transcript.


Background

The product of this gene functions in the NF-kappaB signal transduction pathway. The encoded protein, and the similar and functionally redundant protein MAP3K7IP2/TAB2, forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in response to stimulation with the pro-inflammatory cytokines TNF or IL-1. Subsequent MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-kappaB transcription factor. The human genome contains a related pseudogene. Alternatively spliced transcript variants have been described, but their biological validity has not been determined.

References

Besse, A., et al. J. Biol. Chem. 282(6):3918-3928(2007)
 Kanayama, A., et al. Mol. Cell 15(4):535-548(2004)
 Jin, G., et al. Proc. Natl. Acad. Sci. U.S.A. 101(7):2028-2033(2004)
 Cheung, P.C., et al. Biochem. J. 378 (PT 1), 27-34 (2004) :
 Bouwmeester, T., et al. Nat. Cell Biol. 6(2):97-105(2004)

Images



MAP3K7IP3 Antibody (N-term) (Cat. #AP17071a) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the MAP3K7IP3 antibody detected the MAP3K7IP3 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.