

# TH1L Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17236b

#### **Product Information**

Application WB, E Primary Accession Q8IXH7

Other Accession A5GFY4, NP\_945327.1

Reactivity Human **Predicted** Pig Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB36990 66247 **Calculated MW** 548-576 **Antigen Region** 

#### **Additional Information**

**Gene ID** 51497

Other Names Negative elongation factor C/D, NELF-C/D, TH1-like protein, NELFCD, NELFD,

TH1, TH1L

Target/Specificity This TH1L antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 548-576 amino acids from the

C-terminal region of human TH1L.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** TH1L Antibody (C-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

#### **Protein Information**

Name NELFCD

Synonyms NELFD, TH1, TH1L

**Function** Essential component of the NELF complex, a complex that negatively

regulates the elongation of transcription by RNA polymerase II

(PubMed:<u>12612062</u>). The NELF complex, which acts via an association with the DSIF complex and causes transcriptional pausing, is counteracted by the

P-TEFb kinase complex (PubMed:10199401).

Cellular Location Nucleus.

**Tissue Location** Widely expressed. Expressed in heart, brain, lung, placenta, liver, skeletal and

cardiac muscle, adrenal, thyroid, kidney and pancreas.


## **Background**

The NELF complex of proteins interacts with the DSIF protein complex to repress transcriptional elongation by RNA polymerase II. The protein encoded by this gene is an essential part of the NELF complex. Alternative translation initiation site usage results in the formation of two isoforms with different N-termini.

#### References

Zou, W., et al. Cancer Sci. 101(10):2156-2162(2010) Yang, Y., et al. J. Cell. Biochem. 109(5):1013-1024(2010) Sun, J., et al. J. Biol. Chem. 285(9):6443-6452(2010) Pal, P., et al. Croat. Med. J. 50(4):361-369(2009) Cheng, C., et al. J. Biol. Chem. 284(13):8786-8796(2009)

### **Images**



TH1L Antibody (C-term) (Cat. #AP17236b) western blot analysis in Hela cell line lysates (35ug/lane). This demonstrates the TH1L antibody detected the TH1L protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.