10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

GRIK5 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17247a

Product Information

Application	WB, E
Primary Accession	Q16478
Other Accession	Q63273, Q61626, NP_002079.3
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB37058
Calculated MW	109265
Antigen Region	101-129

Additional Information

Gene ID	2901
Other Names	Glutamate receptor ionotropic, kainate 5, GluK5, Excitatory amino acid receptor 2, EAA2, Glutamate receptor KA-2, KA2, GRIK5, GRIK2
Target/Specificity	This GRIK5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 101-129 amino acids from the N-terminal region of human GRIK5.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GRIK5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GRIK5
Synonyms	GRIK2

Function	Ionotropic glutamate receptor that functions as a cation- permeable ligand-gated ion channel, gated by L-glutamate and the glutamatergic agonist kainic acid. Cannot form functional channels on its own and produces channel activity only in heteromeric assembly with GRIK1 and GRIK2 subunits (PubMed:1321949, PubMed:14511640, PubMed:8730589). Can form functional heteromeric receptors with GRIK3 (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane {ECO:0000250|UniProtKB:Q61626}; Multi-pass membrane protein. Presynaptic cell membrane {ECO:0000250|UniProtKB:Q61626}; Multi-pass membrane protein. Note=Association with GRIK2 is required for its cell membrane localization and channel activity


Background

This gene encodes a protein that belongs to the glutamate-gated ionic channel family. Glutamate functions as the major excitatory neurotransmitter in the central nervous system through activation of ligand-gated ion channels and G protein-coupled membrane receptors. The protein encoded by this gene forms functional heteromeric kainate-preferring ionic channels with the subunits encoded by related gene family members. [provided by RefSeq].

References

Gratacos, M., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 150B (6), 808-816 (2009) :
Gill, M.B., et al. J. Biol. Chem. 284(21):14503-14512(2009)
Arai, S., et al. Psychiatr. Genet. 19(1):6-13(2009)
Fotheringham, J., et al. J Neuroimmune Pharmacol 3(2):105-116(2008)
Shibata, H., et al. Psychiatry Res 141(1):39-51(2006)

Images



GRIK5 Antibody (N-term) (Cat. #AP17247a) western blot analysis in CEM cell line lysates (35ug/lane).This demonstrates the GRIK5 antibody detected the GRIK5 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.