

C9orf7 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP17389b

Product Information

Application	WB, E
Primary Accession	Q9UGQ2
Other Accession	D4A9I3 , Q8BG21 , NP_060056.1 , NP_001129247.1
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB35610
Calculated MW	18470
Antigen Region	123-151

Additional Information

Gene ID	11094
Other Names	Calcium channel flower homolog, Calcium channel flower domain-containing protein 1, CACFD1, C9orf7
Target/Specificity	This C9orf7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 123-151 amino acids from the C-terminal region of human C9orf7.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	C9orf7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CACFD1
Synonyms	C9orf7

Function	Transmembrane protein which mediates synaptic endocytosis and fitness-based cell culling (PubMed: 31341286 , PubMed: 37348560). In response to different stimulus strengths, controls two major modes of synaptic vesicle (SV) retrieval in hippocampal neurons; Clathrin- mediated endocytosis (CME) in response to mild stimulation and activity-dependent bulk endocytosis (ADBE) in response to strong stimulation (By similarity). In cytotoxic T-lymphocytes (CTLs) facilitates calcium-dependent endocytosis of cytotoxic granules at the immuno synapse (By similarity). Different isoforms work as fitness fingerprints in 'loser' and 'winner' cells and thereby mediate win/lose decisions as part of the cell competition process (PubMed: 31341286).
Cellular Location	Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250 UniProtKB:D4A9I3}. Golgi apparatus {ECO:0000250 UniProtKB:D4A9I3}. Vesicle {ECO:0000250 UniProtKB:Q8BG21} Note=In cytotoxic T-lymphocytes, localizes to intracellular vesicles that move to the immuno synapse (By similarity). Enriched in synaptic vesicles at the presynaptic vesicles (By similarity). Detected in the Golgi apparatus of cultured hippocampal neurons (By similarity) {ECO:0000250 UniProtKB:D4A9I3, ECO:0000250 UniProtKB:Q8BG21} [Isoform 2]; Endoplasmic reticulum membrane; Multi-pass membrane protein
Tissue Location	Detected in skin cells at low levels of expression (at protein level).


Background

C9orf7 belongs to the calcium channel flower family.

References

Yao, C.K., et al. Cell 138(5):947-960(2009)
Yuan, X., et al. Am. J. Hum. Genet. 83(4):520-528(2008)
Humphray, S.J., et al. Nature 429(6990):369-374(2004)
Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003)

Images



C9orf7 Antibody (C-term) (Cat. #AP17389b) western blot analysis in mouse stomach tissue lysates (35ug/lane). This demonstrates the C9orf7 antibody detected the C9orf7 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.