

NFIB Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17409c

Product Information

Application WB, E **Primary Accession** 000712

Other Accession P97863, P17924, Q0VCL6, NP 005587.2

Reactivity Human, Mouse

Predicted Bovine, Chicken, Mouse

HostRabbitClonalityPolyclonalIsotypeRabbit IgGClone NamesRB36790Calculated MW47442Antigen Region255-283

Additional Information

Gene ID 4781

Other Names Nuclear factor 1 B-type, NF1-B, Nuclear factor 1/B, CCAAT-box-binding

transcription factor, CTF, Nuclear factor I/B, NF-I/B, NFI-B, TGGCA-binding

protein, NFIB

Target/Specificity This NFIB antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 255-283 amino acids from the Central

region of human NFIB.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions NFIB Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name NFIB

Function Transcriptional activator of GFAP, essential for proper brain development

(PubMed:30388402). Recognizes and binds the palindromic sequence 5'-TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.

Cellular Location

Nucleus.


Background

NFIB recognizes and binds the palindromic sequence 5'-TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.

References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Persson, M., et al. Proc. Natl. Acad. Sci. U.S.A. 106(44):18740-18744(2009)
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Pierron, A., et al. Cancer Genet. Cytogenet. 195(1):66-70(2009)
Le Clerc, S., et al. J. Infect. Dis. 200(8):1194-1201(2009)

Images



Anti-NFIB Antibody (Center) at 1:2000 dilution + human heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 47 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

• MicroRNA 373 Facilitates the Replication of Porcine Reproductive and Respiratory Syndrome Virus by Its Negative Regulation of Type I Interferon Induction.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.