

TOR3A Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17612c

Product Information

Application WB, E Primary Accession Q9H497

Other Accession <u>Q5M936</u>, <u>NP_071766.2</u>

Reactivity Human **Predicted** Rat Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB37650 **Calculated MW** 46199 187-213 **Antigen Region**

Additional Information

Gene ID 64222

Other Names Torsin-3A, ATP-dependent interferon-responsive protein, Torsin family 3

member A, TOR3A, ADIR

Target/Specificity This TOR3A antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 187-213 amino acids from the Central

region of human TOR3A.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

PrecautionsTOR3A Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name TOR3A

Synonyms ADIR

Cellular Location Cytoplasm. Endoplasmic reticulum lumen.

Tissue Location Ubiquitously expressed. Highest expression in stomach, salivary glands and

lymph nodes. Isoform 2 is expressed in placenta.


Background

The function of TOR3A remains unknown.

References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010): Davila, S., et al. Genes Immun. 11(3):232-238(2010) Lamesch, P., et al. Genomics 89(3):307-315(2007) Dron, M., et al. Genomics 79(3):315-325(2002) Ozelius, L.J., et al. Genomics 62(3):377-384(1999)

Images



TOR3A Antibody (Center) (Cat. #AP17612c) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the TOR3A antibody detected the TOR3A protein (arrow).

Citations

• Open-gate mutants of the mammalian proteasome show enhanced ubiquitin-conjugate degradation.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.